



PPAR γ siRNA (h): sc-29455

BACKGROUND

Peroxisome proliferator-activated receptors (PPARs) are nuclear hormone receptors that can be activated by a variety of compounds including fibrates, thiazolidinediones, prostaglandins and fatty acids. Three PPAR subtypes, designated PPAR α , PPAR β (also designated PPAR δ) and PPAR γ , have been described. PPARs promote transcription by forming heterodimers with members of the retinoid X receptor (RXR) family of steroid receptors and binding to specific DNA motifs termed PPAR-response elements (PPREs). PPAR α is abundant in primary hepatocytes where it regulates the expression of proteins involved in fatty acid metabolism. PPAR β is the most widely distributed subtype and is often expressed at high levels. PPAR γ is predominantly seen in adipose tissue where it plays a critical role in regulating adipocyte differentiation. Interestingly, both the orphan nuclear hormone receptor LXR α and thyroid receptor (TR) have been shown to act as antagonists of PPAR α /RXR α binding to PPREs.

REFERENCES

1. Brun, R.P., et al. 1996. Differential activation of adipogenesis by multiple PPAR isoforms. *Genes Dev.* 10: 974-984.
2. Mansen, A., et al. 1996. Expression of the peroxisome proliferator-activated receptor (PPAR) in the mouse colonic mucosa. *Biochem. Biophys. Res. Commun.* 222: 844-851.
3. Sterchele, P.F., et al. 1996. Regulation of peroxisome proliferator-activated receptor- α mRNA in rat liver. *Arch. Biochem. Biophys.* 326: 281-289.
4. Braissant, O., et al. 1996. Differential expression of peroxisome proliferator-activated receptors (PPARs): tissue distribution of PPAR α , β , and γ in the adult rat. *Endocrinology* 137: 354-366.

CHROMOSOMAL LOCATION

Genetic locus: PPARG (human) mapping to 3p25; Pparg (mouse) mapping to 6 E3-F1.

PRODUCT

PPAR γ siRNA (h) is a target-specific 20-25 nt siRNA designed to knock down gene expression. Each vial contains 3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20 $^{\circ}$ C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20 $^{\circ}$ C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

PPAR γ siRNA (h) is recommended for the inhibition of PPAR γ expression in human cells.

PPAR γ (E-8): sc-7273 is recommended as a control antibody for Western Blotting (starting dilution 1:100, dilution range 1:100-1:1,000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) protein detection using the recommended secondary reagents listed below.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-mouse IgG-HRP: sc-2005 (dilution range: 1:2000-1:32,000) or Cruz MarkerTM compatible goat anti-mouse IgG-HRP: sc-2031 (dilution range: 1:2000-1:5000), Cruz MarkerTM Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-mouse IgG-FITC: sc-2010 (dilution range: 1:100-1:400) or goat anti-mouse IgG-TR: sc-2781 (dilution range: 1:100-1:400) with UltraCruzTM Mounting Medium: sc-24941.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 60 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238. Semi-quantitative RT-PCR may be performed using RT-PCR Primer: PPAR γ (h)-PR: sc-29455-PR (20 μ l, 596 bp). Annealing temperature for the primers should be 55-60 $^{\circ}$ C and the extension temperature should be 68-72 $^{\circ}$ C.

SELECT PRODUCT CITATIONS

1. Polikandriotis, J.A., et al. 2005. Peroxisome proliferator-activated receptor gamma ligands stimulate endothelial nitric oxide production through distinct peroxisome proliferator-activated receptor gamma-dependent mechanisms. *Arterioscler. Thromb. Vasc. Biol.* 25: 1810-1816.

RESEARCH USE

For research use only, not for use in diagnostic procedures.