

PPAR β siRNA (h): sc-36305

BACKGROUND

Peroxisome proliferator-activated receptors (PPARs) are nuclear hormone receptors that can be activated by a variety of compounds including fibrates, thiazolidinediones, prostaglandins and fatty acids. Three PPAR subtypes, designated PPAR α , PPAR β (also designated PPAR δ) and PPAR γ , have been described. PPARs promote transcription by forming heterodimers with members of the retinoid X receptor (RXR) family of steroid receptors and binding to specific DNA motifs termed PPAR-response elements (PPREs). PPAR α is abundant in primary hepatocytes where it regulates the expression of proteins involved in fatty acid metabolism. PPAR β is the most widely distributed subtype and is often expressed at high levels. PPAR γ is predominantly seen in adipose tissue where it plays a critical role in regulating adipocyte differentiation. Interestingly, both the orphan nuclear hormone receptor LXR α and thyroid receptor (TR) have been shown to act as antagonists of PPAR α /RXR α binding to PPREs.

REFERENCES

1. Brun, R.P., et al. 1996. Differential activation of adipogenesis by multiple PPAR isoforms. *Genes Dev.* 10: 974-984.
2. Mansen, A., et al. 1996. Expression of the peroxisome proliferator-activated receptor (PPAR) in the mouse colonic mucosa. *Biochem. Biophys. Res. Commun.* 222: 844-851.
3. Lemberger, T., et al. 1996. Expression of the peroxisome proliferator-activated receptor α gene is stimulated by stress and follows a diurnal rhythm. *J. Biol. Chem.* 271: 1764-1769.
4. Braissant, O., et al. 1996. Differential expression of peroxisome proliferator-activated receptors (PPARs): tissue distribution of PPAR- α , - β and - γ in the adult rat. *Endocrinol.* 137: 354-366.

CHROMOSOMAL LOCATION

Genetic locus: PPAR δ (human) mapping to 6p21.1; Ppard (mouse) mapping to 17 A3.3.

PRODUCT

PPAR β siRNA (h) is a pool of 3 target-specific 20-25 nt siRNAs designed to knock down gene expression. Each vial contains 3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections.

For independent verification of PPAR β (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3 nmol of lyophilized siRNA. These include: sc-36305A, sc-36305B and sc-36305C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20 $^{\circ}$ C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20 $^{\circ}$ C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

PPAR β siRNA (h) is recommended for the inhibition of PPAR β expression in human cells.

PPAR β (H-74): sc-7197 is recommended as a control antibody for Western Blotting (starting dilution 1:100, dilution range 1:100-1:1,000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) protein detection using the recommended secondary reagents listed below.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz MarkerTM compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz MarkerTM Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruzTM Mounting Medium: sc-24941.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 60 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238. Semi-quantitative RT-PCR may be performed using RT-PCR Primer: PPAR β (h)-PR: sc-36305-PR (20 μ l, 474 bp). Annealing temperature for the primers should be 55-60 $^{\circ}$ C and the extension temperature should be 68-72 $^{\circ}$ C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.