

Amyloid A4 (I-17): sc-7497

BACKGROUND

Proteolytic cleavage of the amyloid protein precursor (APP) gives rise to the β -amyloid and Amyloid A4 proteins, which are present in human platelets. Amyloid deposition is associated with type II diabetes, Down's syndrome and a variety of neurological disorders, including Alzheimer's disease. The amyloid precursor protein (APP) undergoes alternative splicing, resulting in several isoforms. Proteolytic cleavage of APP leads to the formation of the 4 kDa amyloid β /A4 protein. This protein is involved in the formation of neurofibrillary tangles and plaques that characterize the senile plaques of Alzheimer patients. APLP1 (amyloid precursor-like protein 1) and APLP2 are structurally similar to APP. Human APLP2 is a membrane-bound sperm protein that contains a region highly homologous to the transmembrane-cytoplasmic domains of APP found in brain plaques of Alzheimer disease patients.

REFERENCES

1. Kosik, K.S. 1992 Alzheimer's disease: a cell perspective. *Science* 256: 780-783.
2. Dyrks, T., Dyrks, E., Monning, U., Urmoneit, B., Turner, J., and Beyreuther, K. 1993. Generation of β A4 from the amyloid protein precursor and fragments thereof. *FEBS Letts.* 335: 89-93.
3. Hirai, S. and Okamoto, K. 1993. Amyloid β /A4 peptide associated with Alzheimer's disease and cerebral amyloid angiopathy. *Int. Med.* 32: 923-925.
4. Arendt, T., Holzer, M., Fruth, R., Bruckner, M.K., and Gartner, U. 1995. Paired helical filament-like phosphorylation of tau, deposition of β /A4-amyloid and memory impairment in rat induced by chronic inhibition of phosphatase 1 and 2A. *Neuroscience* 69: 691-698.
5. Gillmore, J.D., Hawkins, P.N., and Pepys, M.B. 1997. Amyloidosis: a review of recent diagnostic and therapeutic developments. *British J. of Haematol.* 99: 245-256.
6. van Leeuwen, F.W., de Kleijn, D.P., van den Hurk, H.H., Neubauer, A., Sonnemans, M.A., Sluijs, J.A., Koycu, S., Ramdjielal, R.D., Salehi, A., Martens, G.J., Grosveld, F.G., Peter, J., Burbach, H., and Hol, E.M. 1998. Frameshift mutants of β amyloid precursor protein and ubiquitin-B in Alzheimer's and Down patients. *Science* 279: 242-247.

SOURCE

Amyloid A4 (I-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of Amyloid A4 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-7497 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

Amyloid A4 (I-17) is recommended for detection of amyloid A4 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for β -Amyloid siRNA (h): sc-29677 and β -Amyloid siRNA (m): sc-29678.


Molecular Weight of Amyloid A4: 100-125 kDa.

Positive Controls: mouse brain extract: sc-2253, rat brain extract: sc-2392 or H4 cell lysate: sc-2408.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Amyloid A4 (I-17): sc-7497. Western blot analysis of Amyloid A4 expression in rat brain (A) and mouse brain (B) tissue extracts.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.