

QTRTD1 (P-14): sc-100112

BACKGROUND

QTRTD1 (queuine tRNA-ribosyltransferase domain containing 1) is a 415 amino acid protein involved in tRNA modification and tRNA-queuosine biosynthesis. Localizing to cytoplasm, QTRTD1 also localizes to the mitochondrial outer membrane and associates with QTRT1 (queuine tRNA-ribosyltransferase domain containing 1) to form an active queuine tRNA-ribosyltransferase. At the wobble position of tRNAs with GUN anticodons, QTRTD1 exchanges queuine for guanine to form queuosine, a modified nucleoside. QTRTD1 is a member of the queuine tRNA-ribosyltransferase family, QTRTD1 subfamily and is encoded by a gene located on human chromosome 3, which houses over 1,100 genes, including a chemokine receptor (CKR) gene cluster and a variety of human cancer-related gene loci. Marfan syndrome, porphyria, von Hippel-Lindau syndrome, osteogenesis imperfecta and Charcot-Marie-Tooth disease are a few of the numerous genetic diseases associated with chromosome 3.

REFERENCES

1. Collod, G., et al. 1994. A second locus for Marfan syndrome maps to chromosome 3p24.2-p25. *Nat. Genet.* 8: 264-268.
2. De Jonghe, P., et al. 1997. Mutilating neuropathic ulcerations in a chromosome 3q13-q22 linked Charcot-Marie-Tooth disease type 2B family. *J. Neurol. Neurosurg. Psychiatr.* 62: 570-573.
3. Maho, A., et al. 1999. Mapping of the CCXCR1, CX3CR1, CCBP2 and CCR9 genes to the CCR cluster within the 3p21.3 region of the human genome. *Cytogenet. Cell Genet.* 87: 265-268.
4. Robinson, P.N., et al. 2000. The molecular genetics of Marfan syndrome and related microfibrilopathies. *J. Med. Genet.* 37: 9-25.
5. Braga, E.A., et al. 2003. New tumor suppressor genes in hot spots of human chromosome 3: new methods of identification. *Mol. Biol.* 37: 194-211.
6. Rasmussen, A., et al. 2010. Uptake of genetic testing and long-term tumor surveillance in von Hippel-Lindau disease. *BMC Med. Genet.* 11: 4.
7. Chen, Y.C., et al. 2010. Characterization of the human tRNA-guanine transglycosylase: confirmation of the heterodimeric subunit structure. *RNA.* 16: 958-968.

CHROMOSOMAL LOCATION

Genetic locus: QTRTD1 (human) mapping to 3q13.31; Qtrtd1 (mouse) mapping to 16 B4.

SOURCE

QTRTD1 (P-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of QTRTD1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-100112 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

QTRTD1 (P-14) is recommended for detection of QTRTD1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for QTRTD1 siRNA (h): sc-78408, QTRTD1 siRNA (m): sc-152617, QTRTD1 shRNA Plasmid (h): sc-78408-SH, QTRTD1 shRNA Plasmid (m): sc-152617-SH, QTRTD1 shRNA (h) Lentiviral Particles: sc-78408-V and QTRTD1 shRNA (m) Lentiviral Particles: sc-152617-V.

Molecular Weight of QTRTD1: 47 kDa.

Positive Controls: HL-60 whole cell lysate: sc-2209, U-937 cell lysate: sc-2239 or LNCaP cell lysate: sc-2231.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.