



UBC13 (YD-16): sc-100622

BACKGROUND

Ubiquitination is an important mechanism through which three classes of enzymes act in concert to target short-lived or abnormal proteins for destruction. The three classes of enzymes involved in ubiquitination are the ubiquitin-activating enzymes (E1s), the ubiquitin-conjugating enzymes (E2s) and the ubiquitin-protein ligases (E3s). UBC13, also known as UBE2N or BLU, is a 152 amino acid member of the E2 ubiquitin-conjugating enzyme family. Existing as a heterodimer with Mms2 (also known as UBE2V2), UBC13 catalyzes the ATP-dependent synthesis of non-canonical polyubiquitin chains, a process that does not lead to proteasomal degradation. Additionally, UBC13 mediates the transcription of several target genes and is thought to play a role in cell cycle progression, cellular differentiation and DNA repair mechanisms that ensure cell survival after DNA damage.

REFERENCES

1. Yamaguchi, T., Kim, N.S., Sekine, S., Seino, H., Osaka, F., Yamao, F. and Kato, S. 1996. Cloning and expression of cDNA encoding a human ubiquitin-conjugating enzyme similar to the *Drosophila* bendless gene product. *J. Biochem.* 120: 494-497.
2. Hoege, C., Pfander, B., Moldovan, G.L., Pyrowolakis, G. and Jentsch, S. 2002. Rad6-dependent DNA repair is linked to modification of PCNA by ubiquitin and SUMO. *Nature* 419: 135-141.
3. Andersen, P.L., Zhou, H., Pastushok, L., Moraes, T., McKenna, S., Ziola, B., Ellison, M.J., Dixit, V.M. and Xiao, W. 2005. Distinct regulation of UBC13 functions by the two ubiquitin-conjugating enzyme variants Mms2 and Uev1A. *J. Cell Biol.* 170: 745-755.
4. Plans, V., Scheper, J., Soler, M., Loukili, N., Okano, Y. and Thomson, T.M. 2006. The RING finger protein RNF8 recruits UBC13 for Lysine 63-based self polyubiquitylation. *J. Cell. Biochem.* 97: 572-582.
5. Yamamoto, M., Okamoto, T., Takeda, K., Sato, S., Sanjo, H., Uematsu, S., Saitoh, T., Yamamoto, N., Sakurai, H., Ishii, K.J., Yamaoka, S., Kawai, T., Matsuura, Y., Takeuchi, O. and Akira, S. 2006. Key function for the UBC13 E2 ubiquitin-conjugating enzyme in immune receptor signaling. *Nat. Immunol.* 7: 962-970.
6. Petroski, M.D., Zhou, X., Dong, G., Daniel-Issakani, S., Payan, D.G. and Huang, J. 2007. Substrate modification with Lysine 63-linked ubiquitin chains through the UBC13-Uev1A ubiquitin-conjugating enzyme. *J. Biol. Chem.* 282: 29936-29945.
7. Zhao, G.Y., Sonoda, E., Barber, L.J., Oka, H., Murakawa, Y., Yamada, K., Ikura, T., Wang, X., Kobayashi, M., Yamamoto, K., Boulton, S.J. and Takeda, S. 2007. A critical role for the ubiquitin-conjugating enzyme UBC13 in initiating homologous recombination. *Mol. Cell* 25: 663-675.
8. Fukushima, T., Matsuzawa, S., Kress, C.L., Bruey, J.M., Krajewska, M., Lefebvre, S., Zapata, J.M., Ronai, Z. and Reed, J.C. 2007. Ubiquitin-conjugating enzyme UBC13 is a critical component of TNF receptor-associated factor (TRAF)-mediated inflammatory responses. *Proc. Natl. Acad. Sci. USA* 104: 6371-6376.

CHROMOSOMAL LOCATION

Genetic locus: UBE2N (human) mapping to 12q22.

SOURCE

UBC13 (YD-16) is a mouse monoclonal antibody raised against recombinant UBC13 of human origin.

PRODUCT

Each vial contains 100 µg IgG_{2b} in 1.0 ml PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

UBC13 (YD-16) is recommended for detection of UBC13 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for UBC13 siRNA (h): sc-43551, UBC13 shRNA Plasmid (h): sc-43551-SH and UBC13 shRNA (h) Lentiviral Particles: sc-43551-V.

Molecular Weight of UBC13: 17 kDa.

Positive Controls: BJAB whole cell lysate: sc-2207 or HeLa whole cell lysate: sc-2200.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-mouse IgG-HRP: sc-2005 (dilution range: 1:2000-1:32,000) or Cruz Marker™ compatible goat anti-mouse IgG-HRP: sc-2031 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.