

B7RP-1 (M-19): sc-10101

BACKGROUND

T-cell co-stimulatory molecule, inducible co-stimulator (ICOS)/B7-related protein-1 (B7RP-1, B7-H2, GL50, ICOS-L) is a ligand for the ICOS receptor that initiates T and B cell proliferation and cytokine secretion. B7RP-1 interactions play an essential role in T cell-dependent B cell activation in peripheral lymphoid organs such as spleen and lymph nodes. B7RP-1 protein is present in myeloid leukocytes, and by northern blot there are 2.4, 3.0, and 7.0 kb transcripts in brain, heart, kidney, and liver, with lower expression in colon and thymus, and a 1.1 kb transcript in leukocytes. Tumor necrosis factor α (TNF α), granulocyte-macrophage colony-stimulating factor (GM-CSF) and interleukin-4 (IL-4) enhance B7RP-1 expression. LPS-induced up-regulation of B7RP-1 is dependent on the MyD88-dependent signaling pathway.

REFERENCES

1. Yoshinaga, S.K., et al. 1999. T cell co-stimulation through B7RP-1 and ICOS. *Nature* 402: 827-832.
2. Ling, V., et al. 2000. Cutting edge: identification of GL50, a novel B7-like protein that functionally binds to ICOS receptor. *J. Immunol.* 164: 1653-1657.
3. Yoshinaga, S.K., et al. 2000. Characterization of a new human B7-related protein: B7RP-1 is the ligand to the co-stimulatory protein ICOS. *Int. Immunol.* 12: 1439-1447.
4. Liyama, R., et al. 2003. The role of inducible co-stimulator (ICOS)/B7-related protein-1 (B7RP-1) interaction in the functional development of Peyer's patches. *Immunol. Lett.* 88: 63-70.
5. Wahl, P., et al. 2003. Interaction of B7RP-1 with ICOS negatively regulates antigen presentation by B cells. *Inflammation* 27: 191-200.

CHROMOSOMAL LOCATION

Genetic locus: Icoslg (mouse) mapping to 10 C1.

SOURCE

B7RP-1 (M-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping within a C-terminal cytoplasmic domain of B7RP-1 of mouse origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-10101 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

B7RP-1 (M-19) is recommended for detection of B7RP-1 of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for B7RP-1 siRNA (m): sc-42769, B7RP-1 shRNA Plasmid (m): sc-42769-SH and B7RP-1 shRNA (m) Lentiviral Particles: sc-42769-V.


Molecular Weight of B7RP-1: 36 kDa.

Positive Controls: B7RP-1 (h): 293T Lysate: sc-116982 or mouse spleen extract: sc-2391.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



B7RP-1 (M-19): sc-10101. Western blot analysis of B7RP-1 expression in non-transfected: sc-117752 (A) and human B7RP-1 transfected: sc-116982 (B) 293T whole cell lysates.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **B7RP-1 (HK5.3): sc-58918**, our highly recommended monoclonal alternative to B7RP-1 (M-19).