

SMC6 (43-Q): sc-101015

BACKGROUND

Breaks in double stranded DNA often arise during DNA replication or as a result of exposure to DNA-damaging agents. Quick and accurate repair of these breaks is crucial for cell survival and genomic stability. Structural maintenance of chromosomes (SMC) family members form heterodimeric complexes that modulate sister chromatid cohesion and chromosome condensation during mitosis. Two distinct SMC protein complexes are the SMC1/SMC3 heterodimer and the SMC2/SMC4 heterodimer. SMC5 and SMC6 play a crucial role in DNA repair as they form a complex that along with SUMO ligase, is also important in preventing DNA damage-induced apoptosis. This complex made up of SMC5 and SMC6 is crucial for sister chromatid homologous recombination DNA repair and also for prevention of chromosomal rearrangements.

REFERENCES

1. Lehmann, A.R. 2005. The role of SMC proteins in the responses to DNA damage. *DNA Repair* 4: 309-314.
2. Lee, K.M. and O'Connell, M.J. 2005. A new SUMO ligase in the DNA damage response. *DNA Repair* 5: 138-141.
3. Potts, P.R. and Yu, H. 2005. Human MMS21/NSE2 is a SUMO ligase required for DNA repair. *Mol. Cell. Biol.* 25: 7021-7032.
4. Watanabe, Y. 2005. The importance of being SMC5/6. *Nat. Cell Biol.* 7: 329-331.
5. Eydmann, T., Machin, F., Dalgaard, J.Z., Farmer, S., Jarmuz, A., Aragon, L. and Torres-Rosell, J. 2005. SMC5 and SMC6 genes are required for the segregation of repetitive chromosome regions. *Nat. Cell Biol.* 7: 412-419.
6. De Piccoli, G., Cortes-Ledesma, F., Ira, G., Torres-Rosell, J., Ceschia, A., Uhle, S., Farmer, S., Hwang, J.Y., Machin, F., McAleenan, A., Papusha, A., Cordon-Preciado, V., Clemente-Blanco, A., Vilella-Mitjana, F., et al. 2006. SMC5-SMC6 mediate DNA double-strand-break repair by promoting sister-chromatid recombination. *Nat. Cell Biol.* 8: 1032-1034
7. Lindroos, H.B., Ström, L., Itoh, T., Katou, Y., Shirahige, K. and Sjögren, C. 2006. Chromosomal association of the SMC5/6 complex reveals that it functions in differently regulated pathways. *Mol. Cell* 22: 755-767.

CHROMOSOMAL LOCATION

Genetic locus: SMC6L1 (human) mapping to 2p24.2.

SOURCE

SMC6 (43-Q) is a mouse monoclonal antibody raised against recombinant SMC6 of human origin.

PRODUCT

Each vial contains 100 µg IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

SMC6 (43-Q) is recommended for detection of SMC6 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for SMC6 siRNA (h): sc-61565, SMC6 shRNA Plasmid (h): sc-61565-SH and SMC6 shRNA (h) Lentiviral Particles: sc-61565-V.

Molecular Weight (predicted) of SMC6 isoforms: 126/129 kDa.


Molecular Weight (observed) of SMC6: 136 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200 or HeLa nuclear extract: sc-2120.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BPHRP: sc-516102 or m-IgGκ BPHRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BPHRP-FITC: sc-516140 or m-IgGκ BPHRP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



SMC6 (43-Q): sc-101015. Western blot analysis of SMC6 expression in HeLa whole cell lysate.



SMC6 (43-Q): sc-101015. Immunoperoxidase staining of formalin-fixed, paraffin-embedded human placenta tissue showing nuclear localization.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.