



TIMP-3 (MM0036-7D3): sc-101578

BACKGROUND

TIMP-1, TIMP-2, TIMP-3 and TIMP-4 (for tissue inhibitor of metalloproteinases 1, 2, 3 and 4) complex with metalloproteinases such as collagenases, gelatinases and stromelysins, resulting in irreversible inactivation of the metalloproteinase. TIMP-1 has been found to be identical to EPA (erythroid-potentiating activity). Parathyroid hormone has been shown to be a regulator of TIMP-2 in osteoblastic cells. TIMP-3 may be involved in regulating trophoblastic invasion of the uterus and remodeling of the extracellular matrix during the folding of epithelia, and in the formation, branching and expansion of epithelial tubes. TIMP-4 is most highly expressed in heart, with low levels expressed in liver, brain, lung, thymus and spleen.

REFERENCES

1. Docherty, A.J., et al. 1985. Sequence of human tissue inhibitor of metalloproteinases and its identity to erythroid-potentiating activity. *Nature* 318: 66-69.
2. Carmichael, D.F., et al. 1986. Primary structure and cDNA cloning of human fibroblast collagenase inhibitor. *Proc. Natl. Acad. Sci. USA* 83: 2407-2411.
3. Cook, T.F., et al. 1994. Cloning and regulation of rat tissue inhibitor of metalloproteinase-2 in osteoblastic cells. *Arch. Biochem. Biophys.* 311: 313-320.
4. Silbiger, S.M., et al. 1994. Cloning of cDNAs encoding human TIMP-3, a novel member of the tissue inhibitor of metalloproteinase family. *Gene* 141: 293-297.
5. Apte, S.S., et al. 1994. Gene encoding a novel murine tissue inhibitor of metalloproteinases (TIMP), TIMP-3, is expressed in developing mouse epithelia, cartilage and muscle, and is located on mouse chromosome 10. *Dev. Dyn.* 200: 177-197.
6. Apte, S.S., et al. 1995. The gene structure of tissue inhibitor of metalloproteinases (TIMP)-3 and its inhibitory activities define the distinct TIMP gene family. *J. Biol. Chem.* 270: 14313-14318.
7. Greene, J., et al. 1996. Molecular cloning and characterization of human tissue inhibitor of metalloproteinase 4. *J. Biol. Chem.* 271: 30375-30380.
8. Gomez, D.E., et al. 1997. Tissue inhibitors of metalloproteinases: structure, regulation and biological functions. *Eur. J. Cell Biol.* 74: 111-122.

CHROMOSOMAL LOCATION

Genetic locus: TIMP3 (human) mapping to 22q12.3.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

SOURCE

TIMP-3 (MM0036-7D3) is a mouse monoclonal antibody raised against recombinant TIMP-3 of human origin.

PRODUCT

Each vial contains 100 µg IgG₁ in 1.0 ml PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

TIMP-3 (MM0036-7D3) is recommended for detection of TIMP-3 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500); non cross-reactive with TIMP-1, TIMP-2 or TIMP-4.

Suitable for use as control antibody for TIMP-3 siRNA (h): sc-37022, TIMP-3 siRNA (h2): sc-44331, TIMP-3 shRNA Plasmid (h): sc-37022-SH, TIMP-3 shRNA Plasmid (h2): sc-44331-SH, TIMP-3 shRNA (h) Lentiviral Particles: sc-37022-V and TIMP-3 shRNA (h2) Lentiviral Particles: sc-44331-V.

Molecular Weight of TIMP-3: 30 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-mouse IgG-HRP: sc-2005 (dilution range: 1:2000-1:32,000) or Cruz Marker™ compatible goat anti-mouse IgG-HRP: sc-2031 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-mouse IgG-FITC: sc-2010 (dilution range: 1:100-1:400) or goat anti-mouse IgG-TR: sc-2781 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 3) Immunohistochemistry: use ImmunoCruz™: sc-2050 or ABC: sc-2017 mouse IgG Staining Systems.