

p-AMPK α 1 (Ser 496): sc-101631

BACKGROUND

AMPK (for 5'-AMP-activated protein kinase) is a heterotrimeric complex comprising a catalytic α subunit and regulatory β and γ subunits. It protects cells from stresses that cause ATP depletion by switching off ATP-consuming biosynthetic pathways. AMPK is activated by high AMP and low ATP through a mechanism involving allosteric regulation, promotion of phosphorylation by an upstream protein kinase known as AMPK kinase, and inhibition of dephosphorylation. Activated AMPK can phosphorylate and regulate *in vivo* hydroxymethylglutaryl-CoA reductase and acetyl-CoA carboxylase, which are key regulatory enzymes of sterol synthesis and fatty acid synthesis, respectively. The human AMPK α 1 and AMPK α 2 genes encode 548 amino acid and 552 amino acid proteins, respectively. Human AMPK β 1 encodes a 271 amino acid protein and human AMPK β 2 encodes a 272 amino acid protein. The human AMPK γ 1 gene encodes a 331 amino acid protein. Human AMPK γ 2 and AMPK γ 3, which are 569 and 492 amino acid proteins, respectively, contain unique N-terminal domains and may participate directly in the binding of AMP within the AMPK complex.

REFERENCES

1. Stapleton, D., et al. 1996. Mammalian AMP-activated protein kinase subfamily. *J. Biol. Chem.* 271: 611-614.
2. Stapleton, D., et al. 1997. AMP-activated protein kinase isoenzyme family: subunit structure and chromosomal location. *FEBS Lett.* 409: 452-456.
3. Hardie, D.G., et al. 1997. The AMP-activated protein kinase-fuel gauge of the mammalian cell? *Eur. J. Biochem.* 246: 259-273.
4. Thornton, C., et al. 1998. Identification of a novel AMP-activated protein kinase γ subunit isoform that is highly expressed in skeletal muscle. *J. Biol. Chem.* 273: 12443-12450.
5. Online Mendelian Inheritance in Man, OMIM[™]. 1998. Johns Hopkins University, Baltimore, MD. MIM Number: 602739. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Cheung, P.C., et al. 2000. Characterization of AMP-activated protein kinase γ subunit isoforms and their role in AMP binding. *Biochem. J.* 346: 659-669.

CHROMOSOMAL LOCATION

Genetic locus: PRKAA1 (human) mapping to 5p13.1; Prkaa1 (mouse) mapping to 15 A1.

SOURCE

p-AMPK α 1 (Ser 496) is a rabbit polyclonal antibody raised against a short amino acid sequence containing phosphorylated Ser 496 of AMPK α 1 of human origin.

PRODUCT

Each vial contains 100 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

p-AMPK α 1 (Ser 496) is recommended for detection of Ser 496 dually phosphorylated AMPK α 1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).


Molecular Weight of p-AMPK α 1: 63 kDa.

Positive Controls: HeLa + heat shock cell lysate: sc-2272 or human colon carcinoma tissue.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto B Blocking Reagent: sc-2335 (use 50 mM NaF, sc-24988, as diluent) and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz[™]: sc-2051 or ABC: sc-2018 rabbit IgG Staining Systems.

DATA



p-AMPK α 1 (Ser 496): sc-101631. Western blot 12 expression in untreated (A) and heat shock-treated (B) HeLa whole cell lysates.



p-AMPK α 1 (Ser 496): sc-101631. Immunoperoxidase staining of formalin-fixed, paraffin-embedded human colon carcinoma tissue showing cytoplasmic localization.

SELECT PRODUCT CITATIONS

1. Sakar, Y., et al. 2010. Metformin-induced regulation of the intestinal D-glucose transporters. *J. Physiol. Pharmacol.* 61: 301-307.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.