

p-EGFR (Ser 1046): sc-101665

BACKGROUND

Epidermal growth factors mediate their effects on cell growth through interactions with a cell surface glycoprotein designated EGFR (EGF receptor). Binding of EGF or TGF α to EGFR activates tyrosine-specific protein kinase activity intrinsic to EGFR. The carboxy terminal tyrosine residues on EGFR, Tyr 1092 and Tyr 1173, designated Tyr 1196 in rat, are the major sites of autophosphorylation which occurs as a result of EGF binding. Once activated, EGFR mediates the binding of the phosphotyrosine binding (PTB) domain of GRB2 through direct interactions with Tyr 1092 and Tyr 1110 in human and mouse or Tyr 1109 in rat, and through indirect interactions with Tyr 1173 in the Ras signaling pathway. Tyr 1173 of EGFR also functions as a kinase substrate. Phosphorylation of Tyr 992, Tyr 1092 and Tyr 1110 is required for conformational change in the C-terminal tail of EGFR. Tyr 1092, Tyr 1173 and Tyr 1110 are also designated Tyr 1068, Tyr 1197, and Tyr 1086, respectively. EGFR is also subject to phosphorylation on other amino acid residues, such as Ser 1046.

REFERENCES

1. Reynolds, F.H. Jr., Todaro, G.J., Fryling, C. and Stephenson, J.R. 1981. Human transforming growth factors induces tyrosine phosphorylation of EGF receptors. *Nature* 292: 259-262.
2. Hunter, T. 1984. The epidermal growth factor receptor gene and its product. *Nature* 311: 414-416.
3. Batzer, A.G., Rotin, D., Urena, J.M., Skolnik, E.Y. and Schlessinger, J. 1994. Hierarchy of binding site for GRB2 and Shc on the epidermal growth factor receptor. *Mol. Cell. Biol.* 14: 5192-5201.
4. Ward, C.W., Gough, K.H., Rashke, M., Wan, S.S., Tribbick, G. and Wang, J-X. 1996. Systematic mapping of potential binding sites for Shc and GRB2 SH2 domains on Insulin receptor substrate-1 and the receptors for Insulin, epidermal growth factor, platelet-derived growth factor, and fibroblast growth factor. *J. Biol. Chem.* 271: 5603-5609.

CHROMOSOMAL LOCATION

Genetic locus: EGFR (human) mapping to 7p11.2; Egfr (mouse) mapping to 11 A2.

SOURCE

p-EGFR (Ser 1046) is a rabbit polyclonal antibody raised against a short amino acid sequence containing phosphorylated Ser 1046 of EGFR of human origin.

PRODUCT

Each vial contains 100 μ g IgG in 1.0 ml PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4 $^{\circ}$ C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

p-EGFR (Ser 1046) is recommended for detection of Ser 1046 phosphorylated EGFR of human and mouse origin and correspondingly phosphorylated Ser 1045 of rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for EGFR siRNA (h): sc-29301, EGFR siRNA (m): sc-29302, EGFR siRNA (r): sc-108050, EGFR shRNA Plasmid (h): sc-29301-SH, EGFR shRNA Plasmid (m): sc-29302-SH, EGFR shRNA Plasmid (r): sc-108050-SH, EGFR shRNA (h) Lentiviral Particles: sc-29301-V, EGFR shRNA (m) Lentiviral Particles: sc-29302-V and EGFR shRNA (r) Lentiviral Particles: sc-108050-V.


Molecular Weight of p-EGFR: 170 kDa.

Positive Controls: A-431 whole cell lysate: sc-2201, Hep G2 cell lysate: sc-2227 or A549 cell lysate: sc-2413.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz MarkerTM compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz MarkerTM Molecular Weight Standards: sc-2035, TBS Blotto B Blocking Reagent: sc-2335 (use 50 mM NaF, sc-24988, as diluent), Western Blotting Luminol Reagent: sc-2048 and Lambda Phosphatase: sc-200312A. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruzTM Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruzTM: sc-2051 or ABC: sc-2018 rabbit IgG Staining Systems.

DATA



p-EGFR (Ser 1070): sc-101665. Western blot analysis of EGFR phosphorylation in untreated (A) and lambda protein phosphatase treated (B) A-431 whole cell lysates.



p-EGFR (Ser 1070): sc-101665. Western blot analysis of phosphorylated EGFR expression in untreated (A) and EGF-treated (B) SK-OV3 whole cell lysates.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.