

# p-c-Jun (Thr 91): sc-101721

## BACKGROUND

Genes belonging to the Jun and Fos oncogene families encode nuclear proteins that are found to be associated with a number of transcriptional complexes. The c-Jun protein is a major component of the transcription factor AP-1, originally shown to mediate phorbol ester tumor promoter (TPA)-induced expression of responsive genes through the TPA-response element (TRE). The Jun proteins form homo- and heterodimers which bind the TRE, while Fos proteins are active only as heterodimers with any of the Jun proteins. Fos/Jun heterodimers have a much higher affinity for the TRE than Jun homodimers. Ha-Ras augments c-Jun activity and stimulates phosphorylation of its activation domain. An inhibitor of Fos/Jun function, termed IP-1, associates with Fos and Jun and is inactivated upon phosphorylation induced by the cAMP-dependent protein kinase A (PKA).

## REFERENCES

1. Sambucetti, L.C., et al. 1986. The Fos protein complex is associated with DNA in isolated nuclei and binds to DNA cellulose. *Science* 234: 1417-1419.
2. Bohmann, D., et al. 1987. Human proto-oncogene c-Jun encodes a DNA binding protein with structural and functional properties of transcription factor AP-1. *Science* 238: 1386-1392.

## CHROMOSOMAL LOCATION

Genetic locus: JUN (human) mapping to 1p32.1; Jun (mouse) mapping to 4 C5.

## SOURCE

p-c-Jun (Thr 91) is a rabbit polyclonal antibody raised against a short amino acid sequence containing Thr 91 phosphorylated c-Jun of human origin.

## PRODUCT

Each vial contains 100 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## APPLICATIONS

p-c-Jun (Thr 91) is recommended for detection of Thr 91 phosphorylated c-Jun of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for c-Jun siRNA (h): sc-29223, c-Jun siRNA (m): sc-29224, c-Jun shRNA Plasmid (h): sc-29223-SH, c-Jun shRNA Plasmid (m): sc-29224-SH, c-Jun shRNA (h) Lentiviral Particles: sc-29223-V and c-Jun shRNA (m) Lentiviral Particles: sc-29224-V.


Molecular Weight of p-c-Jun: 39 kDa.

Positive Controls: A-431 nuclear extract: sc-2122, HeLa nuclear extract: sc-2120 or HeLa + UV cell lysate: sc-2221.


## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto B Blocking Reagent: sc-2335 (use 50 mM NaF, sc-24988, as diluent), Western Blotting Luminol Reagent: sc-2048 and Lambda Phosphatase: sc-200312A. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2051 or ABC: sc-2018 rabbit IgG Staining Systems.

## DATA



p-c-Jun (Thr 91): sc-101721. Western blot analysis of phosphorylated c-Jun expression in untreated (A) and UV-treated (B) HeLa whole cell lysates.



p-c-Jun (Thr 91): sc-101721. Immunoperoxidase staining of formalin-fixed, paraffin-embedded human breast carcinoma tissue showing nuclear localization.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.


 MONOS  
Satisfaction  
Guaranteed

Try **p-c-Jun (C-J 4C4/1): sc-53182**, our highly recommended monoclonal alternative to p-c-Jun (Thr 91).