

p-NFκB p65 (Ser 468): sc-101750

BACKGROUND

Proteins encoded by the v-Rel viral oncogene and its cellular homolog, c-Rel, are members of a family of transcription factors that include the two subunits of the transcription factor NFκB (p50 and p65) and the *Drosophila* maternal morphogen, Dorsal. Both proteins specifically bind to DNA sequences that are the same or slight variations of the 10 bp kB sequence in the immunoglobulin κ light chain enhancer. This same sequence is also present in a number of other cellular and viral enhancers. The DNA binding activity of NFκB is activated and NFκB is subsequently transported from the cytoplasm to the nucleus in cells exposed to mitogens or growth factors. cDNAs encoding precursors for two distinct proteins have been described, designated p105 and p100. The p105 precursor contains p50 at its N-terminus and a C-terminal region that when expressed as a separate molecule, designated PDI, binds to p50 and regulates its activity. The NFκB transcription factor is a protein complex consisting of a DNA binding subunit and an associated protein. The DNA binding subunit, also referred to as Rel A, is functionally related to c-Rel p75 and RelB p68. NFκB p65 is phosphorylated at Serine 311 as a response to protein kinase C ζ

REFERENCES

1. Meyer, R., et al. 1991. Cloning of the DNA binding subunit of human NFκB: the level of its mRNA is strongly regulated by phorbol ester or tumor necrosis factor α. *Proc. Natl. Acad. Sci. USA* 88: 966-970.
2. Schmid, R.M., et al. 1991. Cloning of an NFκB subunit which stimulates HIV transcription in synergy with p65. *Nature* 352: 733-736.
3. Perkins, N.D., et al. 1992. Distinct combinations of NFκB subunits determine the specificity of transcriptional activation. *Proc. Natl. Acad. Sci. USA* 89: 1529-1533.
4. Ballard, D.W., et al. 1992. The 65 kDa subunit of human NFκB functions as a potent transcriptional activator and a target for v-Rel-mediated repression. *Proc. Natl. Acad. Sci. USA* 89: 1875-1879.
5. Hatada, E.N., et al. 1992. The Ankyrin repeat domains of the NFκB precursor p105 and the proto-oncogene Bcl-3 act as specific inhibitors of NFκB DNA binding. *Proc. Natl. Acad. Sci. USA* 89: 2489-2493.
6. Vermeulen, L., et al. 2003. Transcriptional activation of the NFκB p65 subunit by mitogen- and stress-activated protein kinase-1 (MSK1). *EMBO J.* 22: 1313-1324.

CHROMOSOMAL LOCATION

Genetic locus: RELA (human) mapping to 11q13.1; Rela (mouse) mapping to 19 A.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

SOURCE

p-NFκB p65 (Ser 468) is a rabbit polyclonal antibody raised against a short amino acid sequence containing phosphorylated Ser 468 of NFκB p65 of human origin.

PRODUCT

Each vial contains 100 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

p-NFκB p65 (Ser 468) is recommended for detection of Ser 468 phosphorylated NFκB p65 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)].

Suitable for use as control antibody for NFκB p65 siRNA (h): sc-29410, NFκB p65 siRNA (h2): sc-44212 and NFκB p65 siRNA (m): sc-29411.


Molecular Weight of p-NFκB p65: 65 kDa.

Positive Controls: HeLa + TNFα cell lysate: sc-2228, TNFα-treated MDA-MB-231 whole cell lysate or human breast carcinoma tissue.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto B Blocking Reagent: sc-2335 (use 50 mM NaF, sc-24988, as diluent) and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

DATA



Western blot analysis of phosphorylated NFκB p65 expression in MOLT-4 (A, B) and HeLa (C) whole cell lysates. Blots were probed with p-NFκB p65 (Ser 468): sc-101750 preincubated with cognate phosphorylated peptide (A) and p-NFκB p65 (Ser 468): sc-101750 (B, C).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.