

# p-SH-PTP2 (Tyr 542): sc-101798

## BACKGROUND

The steady state of protein tyrosyl phosphorylation in cells is regulated by the opposing action of tyrosine kinases and protein tyrosine phosphatases (PTPs). Several groups have independently identified a non-transmembrane PTP, designated SH-PTP1 (also known as PTP1C, HCP and SHP), which is primarily expressed in hematopoietic cells and characterized by the presence of two SH2 domains N-terminal to the PTP domain. SH2 domains generally mediate the association of regulatory molecules with specific phosphotyrosine-containing sites on autophosphorylated receptors, thereby controlling the initial interaction of receptors with these substrates. A second and much more widely expressed PTP with SH2 domains, SH-PTP2 (also designated PTP1D and Syp), has been identified. Strong sequence similarity between SH-PTP2 and the *Drosophila* gene corkscrew (CSW) and their similar patterns of expression suggest that SH-PTP2 is the human corkscrew homolog.

## CHROMOSOMAL LOCATION

Genetic locus: PTPN11 (human) mapping to 12q24.13; Ptpn11 (mouse) mapping to 5 F.

## SOURCE

p-SH-PTP2 (Tyr 542) is a rabbit polyclonal antibody raised against a short amino acid sequence containing Tyr 542 phosphorylated SH-PTP2 of human origin.

## PRODUCT

Each vial contains 100 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

p-SH-PTP2 (Tyr 542) is recommended for detection of Tyr 542 phosphorylated SH-PTP2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for SH-PTP2 siRNA (h): sc-36488, SH-PTP2 siRNA (m): sc-36489, SH-PTP2 siRNA (r): sc-270045, SH-PTP2 shRNA Plasmid (h): sc-36488-SH, SH-PTP2 shRNA Plasmid (m): sc-36489-SH, SH-PTP2 shRNA Plasmid (r): sc-270045-SH, SH-PTP2 shRNA (h) Lentiviral Particles: sc-36488-V, SH-PTP2 shRNA (m) Lentiviral Particles: sc-36489-V and SH-PTP2 shRNA (r) Lentiviral Particles: sc-270045-V.


Molecular Weight of p-SH-PTP2: 70 kDa.

Positive Controls: insulin-treated NIH/3T3 whole cell lysate.


## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto B Blocking Reagent: sc-2335 (use 50 mM NaF, sc-24988, as diluent), Western Blotting Luminol Reagent: sc-2048 and Lambda Phosphatase: sc-200312A. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2051 or ABC: sc-2018 rabbit IgG Staining Systems.

## DATA



p-SH-PTP2 (Tyr 542): sc-101798. Western blot analysis of phosphorylated SH-PTP2 expression in untreated (A) and insulin-treated (B) NIH/3T3 whole cell lysates.



p-SH-PTP2 (Tyr 542): sc-101798. Immunoperoxidase staining of formalin-fixed, paraffin-embedded human breast carcinoma tissue showing nuclear localization.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.

**MONOS**  
Satisfaction  
Guaranteed

Try **p-SH-PTP2 (49.Tyr 542): sc-293147**, our highly recommended monoclonal alternative to p-SH-PTP2 (Tyr 542).