

p-SH-PTP2 (Tyr 580): sc-101799

BACKGROUND

The steady state of protein tyrosyl phosphorylation in cells is regulated by the opposing action of tyrosine kinases and protein tyrosine phosphatases (PTPs). Several groups have independently identified a non-transmembrane PTP, designated SH-PTP1 (also known as PTP1C, HCP and SHP), which is primarily expressed in hematopoietic cells and characterized by the presence of two SH2 domains N-terminal to the PTP domain. SH2 domains generally mediate the association of regulatory molecules with specific phosphotyrosine-containing sites on autophosphorylated receptors, thereby controlling the initial interaction of receptors with these substrates. A second and much more widely expressed PTP with SH2 domains, SH-PTP2 (also designated PTP1D and Syp), has been identified. Strong sequence similarity between SH-PTP2 and the *Drosophila* gene corkscrew (CSW) and their similar patterns of expression suggest that SH-PTP2 is the human corkscrew homolog.

REFERENCES

- Chernoff, J., et al. 1990. Cloning of a cDNA for a major human protein-tyrosine-phosphatase. Proc. Natl. Acad. Sci. USA 87: 2735-2739.
- Shen, S., et al. 1991. A protein-tyrosine phosphatase with sequence similarity to the SH2 domain of the protein-tyrosine kinases. Nature 352: 736-739.
- Plutzky, J., et al. 1992. Isolation of a Src homology 2-containing tyrosine phosphatase. Proc. Natl. Acad. Sci. USA 89: 1123-1127.
- Freeman, R.M. Jr., et al. 1992. Identification of a human Src homology 2-containing protein-tyrosine-phosphatase: a putative homolog of *Drosophila* corkscrew. Proc. Natl. Acad. Sci. USA 89: 11239-11243.
- Yi, T., et al. 1992. Protein tyrosine phosphatase containing SH2 domains: characterization, preferential expression in hematopoietic cells and localization to human chromosome 12p12-p13. Mol. Cell. Biol. 12: 836-846.
- Matthews, R.J., et al. 1992. Characterization of hematopoietic intracellular protein tyrosine phosphatases: description of a phosphatase containing an SH2 domain and another enriched in proline-, glutamic acid-, serine- and threonine-rich sequences. Mol. Cell. Biol. 12: 2396-2405.

CHROMOSOMAL LOCATION

Genetic locus: PTPN11 (human) mapping to 12q24.13; Ptpn11 (mouse) mapping to 5 F.

SOURCE

p-SH-PTP2 (Tyr 580) is a rabbit polyclonal antibody raised against a short amino acid sequence containing phosphorylated Tyr 580 of SH-PTP2 of human origin.

PRODUCT

Each vial contains 100 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

p-SH-PTP2 (Tyr 580) is recommended for detection of Tyr 580 phosphorylated SH-PTP2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for SH-PTP2 siRNA (h): sc-36488, SH-PTP2 siRNA (m): sc-36489, SH-PTP2 siRNA (r): sc-270045, SH-PTP2 shRNA Plasmid (h): sc-36488-SH, SH-PTP2 shRNA Plasmid (m): sc-36489-SH, SH-PTP2 shRNA Plasmid (r): sc-270045-SH, SH-PTP2 shRNA (h) Lentiviral Particles: sc-36488-V, SH-PTP2 shRNA (m) Lentiviral Particles: sc-36489-V and SH-PTP2 shRNA (r) Lentiviral Particles: sc-270045-V.


Molecular Weight of p-SH-PTP2: 70 kDa.

Positive Controls: human breast carcinoma tissue.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto B Blocking Reagent: sc-2335 (use 50 mM NaF, sc-24988, as diluent) and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 3) Immunohistochemistry: use ImmunoCruz™: sc-2051 or ABC: sc-2018 rabbit IgG Staining Systems.

DATA



p-SH-PTP2 (Tyr 580): sc-101799. Immunoperoxidase staining of formalin-fixed, paraffin-embedded human breast carcinoma tissue showing cytoplasmic staining.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.