

MOX-2 (C-15): sc-10189

BACKGROUND

Closely related homeobox proteins, MOX-1 and MOX-2, belong to a family of nonclustered, diverged homeobox genes that are expressed in overlapping patterns in the paraxial mesoderm and its derivatives. MOX-1 and MOX-2 function transiently in the formation of mesodermal and mesenchymal derivatives. MOX-1 and MOX-2 are implicated in the early steps of mesoderm formation during gastrulation. In addition, the MOX proteins are also involved in somatic differentiation. Significantly, MOX-1 associates more strongly with Pax1, whereas MOX-2 preferentially associates with Pax3. Specifically, expression of MOX-2, also known as mesenchyme homeobox 2 and GAX, has been shown to be critical in axial skeleton development. MOX-2 is not needed for the migration of myogenic precursors into the limb bud, but it is essential for normal appendicular muscle formation and for the normal regulation of myogenic genes. MOX-2 is expressed in placental tissue. The human MEOX2 gene maps to chromosome 7p21.2 and encodes the MOX-2 protein. Mutations in the gene may be involved in craniofacial and/or skeletal abnormalities.

CHROMOSOMAL LOCATION

Genetic locus: MEOX2 (human) mapping to 7p21.2; Meox2 (mouse) mapping to 12 A3.

SOURCE

MOX-2 (C-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of MOX-2 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-10189 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

MOX-2 (C-15) is recommended for detection of MOX-2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for MOX-2 siRNA (h): sc-106233, MOX-2 siRNA (m): sc-149520, MOX-2 shRNA Plasmid (h): sc-106233-SH, MOX-2 shRNA Plasmid (m): sc-149520-SH, MOX-2 shRNA (h) Lentiviral Particles: sc-106233-V and MOX-2 shRNA (m) Lentiviral Particles: sc-149520-V.


Molecular Weight of MOX-2: 34 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200 or MOX-2 (h): 293 Lysate: sc-113256.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

DATA



MOX-2 (C-15): sc-10189. Western blot analysis of MOX-2 expression in non-transfected: sc-117750 (A) and human MOX-2 transfected: sc-113256 (B) whole cell lysates.



MOX-2 (C-15): sc-10189. Immunoperoxidase staining of formalin fixed, paraffin-embedded human placenta tissue showing nuclear and cytoplasmic staining of decidual cells.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfaction
Guaranteed

Try **MOX-2 (A-8): sc-376748** or **MOX-2 (F-7): sc-390075**, our highly recommended monoclonal alternatives to MOX-2 (C-15).