

# BTBD14A (J-18): sc-101897

## BACKGROUND

BTBD14A (BTB/POZ domain-containing protein 14A), also known as BTBD14, is a 587 amino acid protein that contains one BTB/POZ domain. The BTB/POZ domain mediates homomeric and heteromeric POZ-POZ interactions and is common to transcriptional regulators involved in chromatin modeling. In several BTB/POZ containing proteins, including Bcl-6 and the promyelocytic leukemia zinc-finger (PLZF) oncoprotein, this domain interacts with the SMRT/N-CoR-mSin3A HDAC complex and is directly involved in repressing and silencing gene transcription. When this domain is deleted, as with the oncogenic PLZF-RAR chimera of promyelocytic leukemias, this transcriptional repression is attenuated. This suggests that BTBD14A may play a role in transcription regulation.

## REFERENCES

1. Wong, C.W. and Privalsky, M.L. 1998. Components of the SMRT co-repressor complex exhibit distinctive interactions with the POZ domain oncoproteins PLZF, PLZF-RAR $\alpha$ , and Bcl-6. *J. Biol. Chem.* 273: 27695-27702.
2. David, G., et al. 1998. Histone deacetylase associated with mSin3A mediates repression by the acute promyelocytic leukemia-associated PLZF protein. *Oncogene* 16: 2549-2556.
3. Huynh, K.D. and Bardwell, V.J. 1998. The Bcl-6 POZ domain and other POZ domains interact with the co-repressors N-CoR and SMRT. *Oncogene* 17: 2473-2484.
4. Ahmad, K.F., et al. 1998. Crystal structure of the BTB domain from PLZF. *Proc. Natl. Acad. Sci. USA* 95: 12123-12128.
5. Deltour, S., et al. 1999. Recruitment of SMRT/N-CoR-mSin3A-HDAC-repressing complexes is not a general mechanism for BTB/POZ transcriptional repressors: the case of HIC-1 and  $\gamma$ FBP-B. *Proc. Natl. Acad. Sci. USA* 96: 14831-14836.
6. Melnick, A., et al. 2002. Critical residues within the BTB domain of PLZF and Bcl-6 modulate interaction with co-repressors. *Mol. Cell. Biol.* 22: 1804-1818.
7. Humphray, S.J., et al. 2004. DNA sequence and analysis of human chromosome 9. *Nature* 429: 369-374.

## CHROMOSOMAL LOCATION

Genetic locus: BTBD14A (human) mapping to 9q34.3; Btdb14a (mouse) mapping to 2 A3.

## SOURCE

BTBD14A (J-18) is a purified rabbit polyclonal antibody raised against BTBD14A of human origin.

## PRODUCT

Each vial contains 100  $\mu$ g IgG in 1.0 ml PBS with < 0.1% sodium azide, 0.1% gelatin and < 0.02% sucrose.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

BTBD14A (J-18) is recommended for detection of BTBD14A of mouse, rat, human and canine origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for BTBD14A siRNA (h): sc-92557, BTBD14A siRNA (m): sc-141772, BTBD14A shRNA Plasmid (h): sc-92557-SH, BTBD14A shRNA Plasmid (m): sc-141772-SH, BTBD14A shRNA (h) Lentiviral Particles: sc-92557-V and BTBD14A shRNA (m) Lentiviral Particles: sc-141772-V.


Molecular Weight of BTBD14A: 63 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204, NIH/3T3 whole cell lysate: sc-2210 or K-562 whole cell lysate: sc-2203.


## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2051 or ABC: sc-2018 rabbit IgG Staining Systems.

## DATA



BTBD14A (J-18): sc-101897. Western blot analysis of BTBD14A expression in mouse skin tissue extract (A) and NIH/3T3 (B), Jurkat (C), A2058 (D), HeLa (E) and K-562 (F) whole cell lysates.



BTBD14A (J-18): sc-101897. Immunoperoxidase staining of formalin fixed, paraffin-embedded human kidney tissue showing cytoplasmic staining.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.