

OATP-I (J-16): sc-102045

BACKGROUND

The organic anion transporter family of proteins mediate hepatic uptake of cardiac glycosides. OATP-I (organic anion transporter F), also known as SLC06A1 (solute carrier organic anion transporter family member 6A1), gonad-specific transporter (GST), cancer/testis antigen 48 (CT48) or SLC21A14 (solute carrier family 21 member 19), is a 719 amino acid member of the organic anion transporter protein family. As a multi-pass membrane protein, OATP-I is thought to mediate the Na⁺-independent transport of organic anions. OATP-I is highly expressed in testis, with lower expression found in brain, placenta and spleen. OATP-I is expressed as three isoforms produced by alternative splicing events.

REFERENCES

- Hsiang, B., Zhu, Y., Wang, Z., Wu, Y., Sasseville, V., Yang, W.P. and Kirchgessner, T.G. 1999. A novel human hepatic organic anion transporting poly-peptide (OATP2). *J. Biol. Chem.* 274: 37161-37168.
- Konig, J., Cui, Y., Nies, A.T. and Keppler, D. 2000. Localization and genomic organization of a new hepatocellular organic anion transporting polypeptide. *J. Biol. Chem.* 275: 23161-23168.
- Cai, S.Y., Wang, W., Soroka, C.J., Ballatori, N. and Boyer, J.L. 2002. An evolutionarily ancient Oatp: insights into conserved functional domains of these proteins. *Am. J. Physiol. Gastrointest. Liver Physiol.* 282: 702-710.
- Pizzagalli, F., Hagenbuch, B., Stieger, B., Klenk, U., Folkers, G. and Meier, P.J. 2002. Identification of a novel human organic anion transporting polypeptide as a high affinity thyroxine transporter. *Mol. Endocrinol.* 16: 2283-2296.
- Patel, P., Weerasekera, N., Hitchins, M., Boyd, C.A., Johnston, D.G. and Williamson, C. 2002. Semi quantitative expression analysis of Mdr-3, FIC1, BSEP, Oatp-A, Oatp-C, Oatp-D, Oatp-E and Ntcp gene transcripts in 1st and 3rd trimester human placenta. *Placenta* 24: 39-44.
- Suzuki, T., Onogawa, T., Asano, N., Mizutamari, H., Mikkaichi, T., Tanemoto, M., Abe, M., Satoh, F., Unno, M., Nunoki, K., Suzuki, M., Hishinuma, T., Goto, J., Shimosegawa, T., Matsuno, S., Ito, S. and Abe, T. 2003. Identification and characterization of novel rat and human gonad-specific organic anion transporters. *Mol. Endocrinol.* 17: 1203-1215.

CHROMOSOMAL LOCATION

Genetic locus: SLC06A1 (human) mapping to 5q21.1.

SOURCE

OATP-I (J-16) is a purified rabbit polyclonal antibody raised against OATP-I of human origin.

PRODUCT

Each vial contains 100 µg IgG in 1.0 ml PBS with < 0.1% sodium azide, 0.1% gelatin and < 0.02% sucrose.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

OATP-I (J-16) is recommended for detection of OATP-I of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for OATP-I siRNA (h): sc-91933, OATP-I shRNA Plasmid (h): sc-91933-SH and OATP-I shRNA (h) Lentiviral Particles: sc-91933-V.


Molecular Weight of OATP-I: 79 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

DATA



OATP-I (J-16): sc-102045. Western blot analysis of OATP-I expression in Jurkat whole cell lysate.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.