SANTA CRUZ BIOTECHNOLOGY, INC.

SAP 155 (P-25): sc-102102



BACKGROUND

SAP 155 (spliceosome-associated protein 155), also known as SF3B1, SF3B155 (splicing factor 3β , subunit 1, 155 kDa), PRP10 or PRPF10, is a 1,304 amino acid member of the SF3B1 family and contains eleven HEAT repeats. Localized to nuclear speckles and also to the cytoplasm during mitosis, SAP 155 is a subunit of the SF3B splicing factor. The SF3B splicing factor is a U2 snRNP-associated protein complex essential for spliceosome assembly. SF3B contains the spliceosomal proteins SAP 49, SAP 130, SAP 145 and SAP 155. Concomitant with splicing catalysis, SAP 155 is phosphorylated at its N-terminal Thr-Pro dipeptide motifs by Dyrk1A and cyclin E/Cdk2. This modification of SAP 155 is vital for a functional spliceosome as it is an essential event in the basic splicing reaction. Due to alternative splicing events, various SAP 155 isoforms are produced.

REFERENCES

- 1. Wang, C., et al. 1998. Phosphorylation of spliceosomal protein SAP 155 coupled with splicing catalysis. Genes Dev. 12: 1409-1414.
- Isono, K., et al. 2001. Molecular cloning, genetic mapping, and expression of the mouse Sf3b1 (SAP 155) gene for the U2 snRNP component of spliceosome. Mamm. Genome 12: 192-198.

CHROMOSOMAL LOCATION

Genetic locus: SF3B1 (human) mapping to 2q33.1; Sf3b1 (mouse) mapping to 1 C1.2.

SOURCE

SAP 155 (P-25) is a purified rabbit polyclonal antibody raised against SAP 155 of human origin.

PRODUCT

Each vial contains 100 μg lgG in 1.0 ml PBS with < 0.1% sodium azide, 0.1% gelatin and < 0.02% sucrose.

APPLICATIONS

SAP 155 (P-25) is recommended for detection of SAP 155 of mouse, rat, human and canine origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for SAP 155 siRNA (h): sc-94471, SAP 155 siRNA (m): sc-153216, SAP 155 shRNA Plasmid (h): sc-94471-SH, SAP 155 shRNA Plasmid (m): sc-153216-SH, SAP 155 shRNA (h) Lentiviral Particles: sc-94471-V and SAP 155 shRNA (m) Lentiviral Particles: sc-153216-V.

Molecular Weight of SAP 155: 155 kDa.

Positive Controls: SAP 155 (m): 293T Lysate: sc-110269, Jurkat nuclear extract: sc-2132 or K-562 nuclear extract: sc-2130.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941. 4) Immuno-histochemistry: use ImmunoCruz[™]: sc-2051 or ABC: sc-2018 rabbit IgG Staining Systems.

DATA





SAP 155 (P-25): sc-102102. Western blot analysis of SAP 155 expression in non-transfected: sc-117752 (**A**) and mouse SAP 155 transfected: sc-110269 (**B**) 2931 whole cell lysates and Jurkat (**C**), K-562 (**D**), Ramos (**E**) and MCF7 (**F**) nuclear extracts.

SAP 155 (P-25): sc-102102. Immunoperoxidase staining of formalin fixed, paraffin-embedded human liver tissue showing nuclear and cytoplasmic staining.

STORAGE

Store at 4° C, **D0 NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS Satisfation Guaranteed

Try SAP 155 (B-3): sc-514655 or SAP 155 (G-9): sc-514494, our highly recommended monoclonal aternatives to SAP 155 (P-25).