

ZNF385A (Q-23): sc-102213

BACKGROUND

Zinc finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. ZNF385A (zinc finger matrin-type protein 385A), also known as HZF (hematopoietic zinc finger protein), RZF (retinal zinc finger protein) or ZNF385, is a 366 amino acid protein that contains three matrin-type zinc fingers. The matrin-type zinc finger, which is very similar in structure to the classical DNA-binding C₂H₂ zinc finger, was first identified in the protein matrin-3. The matrin-type zinc finger has also been identified in several spliceosome RNA-binding proteins, suggesting a role in pre-mRNA binding. ZNF385A is expressed predominantly in the retina, localized to the nucleus as well as the cytoplasm. Two isoforms of ZNF385A exist due to alternative splicing events.

REFERENCES

- Thiesen, H.J. 1990. Multiple genes encoding zinc finger domains are expressed in human T cells. *New Biol.* 2: 363-374.
- Belgrader, P., Dey, R. and Berezney, R. 1991. Molecular cloning of matrin-3. A 125-kilodalton protein of the nuclear matrix contains an extensive acidic domain. *J. Biol. Chem.* 266: 9893-9899.
- Rossi, F., Forné, T., Antoine, E., Tazi, J., Brunel, C. and Cathala, G. 1996. Involvement of U1 small nuclear ribonucleoproteins (SnRNP) in 5' splice site-U1 SnRNP interaction. *J. Biol. Chem.* 271: 23985-23991.
- Matsushima, Y., Matsumura, K. and Kitagawa, Y. 1997. Zinc finger-like motif conserved in a family of RNA binding proteins. *Biosci. Biotechnol. Biochem.* 61: 905-906.
- Hibino, Y. 2000. Functional arrangement of genomic DNA and structure of nuclear matrix. *Yakugaku Zasshi* 120: 520-533.
- Durand, S., Abadie, P., Angeletti, S. and Genti-Raimondi, S. 2003. Identification of multiple differentially expressed messenger RNAs in normal and pathological trophoblast. *Placenta* 24: 209-218.
- Muto, Y., Pomeranz Krummel, D., Oubridge, C., Hernandez, H., Robinson, C.V., Neuhaus, D. and Nagai, K. 2004. The structure and biochemical properties of the human spliceosomal protein U1C. *J. Mol. Biol.* 341: 185-198.
- Ainscough, J.F., Rahman, F.A., Sercombe, H., Sedo, A., Gerlach, B. and Coverley, D. 2007. C-terminal domains deliver the DNA replication factor CIZ1 to the nuclear matrix. *J. Cell Sci.* 120 Pt. 1: 115-124.

CHROMOSOMAL LOCATION

Genetic locus: ZNF385A (human) mapping to 12q13.13; Zfp385a (mouse) mapping to 15 F3.

SOURCE

ZNF385A (Q-23) is a purified rabbit polyclonal antibody raised against ZNF385A of human origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 50 µg IgG in 500 µl PBS with < 0.1% sodium azide, 0.1% gelatin and < 0.02% sucrose.

APPLICATIONS

ZNF385A (Q-23) is recommended for detection of ZNF385A of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for ZNF385A siRNA (h): sc-95676, ZNF385A siRNA (m): sc-155704, ZNF385A shRNA Plasmid (h): sc-95676-SH, ZNF385A shRNA Plasmid (m): sc-155704-SH, ZNF385A shRNA (h) Lentiviral Particles: sc-95676-V and ZNF385A shRNA (m) Lentiviral Particles: sc-155704-V.


Molecular Weight of ZNF385A: 38 kDa.

Positive Controls: ZNF385 (h): 293T Lysate: sc-113745.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2051 or ABC: sc-2018 rabbit IgG Staining Systems.

DATA



ZNF385 (Q-23): sc-102213. Western blot analysis of ZNF385 expression in non-transfected 293T: sc-117752 (A), human ZNF385 transfected 293T: sc-113745 (B) and HeLa (C) whole cell lysates.



ZNF385 (Q-23): sc-102213. Immunoperoxidase staining of formalin fixed, paraffin-embedded human lung tissue showing nuclear and cytoplasmic staining.

RESEARCH USE

For research use only, not for use in diagnostic procedures.