


9G8 (G-14): sc-10245



The Power to Question

BACKGROUND

Pre-mRNA splicing enhancer elements are short RNA sequences capable of activating weak splice sites in nearby introns, and they are required for accurate splice site recognition and the control of alternative splicing. Splicing enhancer elements contain specific binding sites for serine/arginine (SR)-rich splicing factors, which include SC35, 9G8, SRp20, and SF2/ASF. The family of SR factors all contain one or more RNA recognition motifs (RRM) and an arginine/serine (RS)-rich domain, and they are essential for constitutive splicing and also regulate splicing in a concentration-dependent manner by influencing the selection of alternative splice sites. The majority of SR proteins, including SC35 and SRp40, are confined to the nucleus, while SF2/ASF, SRp20, and 9G8 are continuously shuttled between the nucleus and the cytoplasm and contribute to mRNA transport. The activity of SR proteins in regulated splicing is antagonized by members of the hnRNP A/B family of proteins, which induce drastic shifts in the selection of splicing-sites. An additional SR-associated protein, p32, tightly associates with SR factors and preferentially inhibits ASF/SF2 functioning as both a splicing enhancer and splicing repressor protein by preventing the stable interaction of ASF/SF2 and the RNA.

REFERENCES

1. Fu, X.D. 1993. Specific commitment of different pre-mRNAs to splicing by single SR proteins. *Nature* 365: 82-85.
2. Mayeda, A., et al. 1994. Function of conserved domains of hnRNP A1 and other hnRNP A/B proteins. *EMBO J.* 13: 5483-5495.
3. Jumaa, H., et al. 1997. The splicing factor SRp20 modifies splicing of its own mRNA and ASF/SF2 antagonizes this regulation. *EMBO J.* 16: 5077-5085.
4. Caceres, J.F., et al. 1998. A specific subset of SR proteins shuttles continuously between the nucleus and the cytoplasm. *Genes Dev.* 12: 55-66.
5. Cavaloc, Y., et al. 1999. The splicing factors 9G8 and SRp20 transactivate splicing through different and specific enhancers. *RNA* 5: 468-483.

CHROMOSOMAL LOCATION

Genetic locus: SFRS7 (human) mapping to 2p22.1.

SOURCE

9G8 (G-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of 9G8 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-10245 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

9G8 (G-14) is recommended for detection of 9G8 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for 9G8 siRNA (h): sc-38246, 9G8 shRNA Plasmid (h): sc-38246-SH and 9G8 shRNA (h) Lentiviral Particles: sc-38246-V.


Molecular Weight of 9G8: 35 kDa.

Positive Controls: Hs 732.Sk/Mu whole cell lysate: sc-364362.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

DATA



9G8 (G-14): sc-10245. Western blot analysis of 9G8 expression in Hs 732.Sk/Mu whole cell lysate.



9G8 (G-14): sc-10245. Immunoperoxidase staining of formalin fixed, paraffin-embedded human cerebral cortex tissue showing nuclear and cytoplasmic staining of neuronal cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **9G8/SRp20 (G-3): sc-390126**, our highly recommended monoclonal alternative to 9G8 (G-14).