



CTDSPL (P-12): sc-102464

BACKGROUND

The family of small CTD phosphatases preferentially catalyze the dephosphorylation of Ser 5 within a consensus repeat. As a member of this family, CTDSPL (CTD small phosphatase-like protein), also known as nuclear LIM interactor-interacting factor 1, RBSP3 or protein YA22, is a 276 amino acid nuclear protein that is ubiquitously expressed. There are two isoforms of CTDSPL that exist as a result of alternative splicing events. Since both of these isoforms reduce the level of phosphorylated Rb protein when transfected into mammary carcinoma cells, CTDSPL functions primarily as a phosphatase. The gene encoding CTDSPL is deleted in 15% of major epithelial cancers, suggesting that CTDSPL plays a role in tumor suppression. Due to evidence showing that inactive small CTD phosphatases interfere with the repressor element 1-silencing transcription factor/neuron-restrictive silencer factor (REST/NRSF) complex, therefore promoting neuronal differentiation, it is suspected that CTDSPL may function as a transcription regulator that acts to silence neuronal genes.

REFERENCES

- Ishikawa, S., Kai, M., Tamari, M., Takei, Y., Takeuchi, K., Bandou, H., Yamane, Y., Ogawa, M. and Nakamura, Y. 1997. Sequence analysis of a 685-kb genomic region on chromosome 3p22-p21.3 that is homozygously deleted in a lung carcinoma cell line. *DNA Res.* 4: 35-43.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 608592. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Yeo, M., Lin, P.S., Dahmus, M.E. and Gill, G.N. 2003. A novel RNA polymerase II C-terminal domain phosphatase that preferentially dephosphorylates Serine 5. *J. Biol. Chem.* 278: 26078-26085.
- Kashuba, V.I., Li, J., Wang, F., Senchenko, V.N., Protopopov, A., Malyukova, A., Kutsenko, A.S., Kadyrova, E., Zabarovska, V.I., Muravenko, O.V., Zelenin, A.V., Kisselev, L.L., Kuzmin, I., Minna, J.D., Winberg, G., Ernberg, I., Braga, E., Lerman, M.I., Klein, G. and Zabarovsky, E.R. 2004. RBSP3 (HYA22) is a tumor suppressor gene implicated in major epithelial malignancies. *Proc. Natl. Acad. Sci. USA* 101: 4906-4911.
- Yeo, M., Lee, S.K., Lee, B., Ruiz, E.C., Pfaff, S.L. and Gill, G.N. 2005. Small CTD phosphatases function in silencing neuronal gene expression. *Science*. 307: 596-600.
- Murabito, J.M., Rosenberg, C.L., Finger, D., Kreger, B.E., Levy, D., Splansky, G.L., Antman, K. and Hwang, S.J. 2007. A genome-wide association study of breast and prostate cancer in the NHLBI's Framingham Heart Study. *BMC Med. Genet.* 8 Suppl1: S6.
- Yeo, M. and Lin, P.S. 2007. Functional characterization of small CTD phosphatases. *Methods Mol. Biol.* 365: 335-346.
- Chang, T.C., Yu, D., Lee, Y.S., Wentzel, E.A., Arking, D.E., West, K.M., Dang, C.V., Thomas-Tikhonenko, A. and Mendell, J.T. 2008. Widespread microRNA repression by Myc contributes to tumorigenesis. *Nat. Genet.* 40: 43-50.

CHROMOSOMAL LOCATION

Genetic locus: CTDSPL (human) mapping to 3p22.2.

SOURCE

CTDSPL (P-12) is an affinity purified rabbit polyclonal antibody raised against a peptide mapping within an internal region of CTDSPL of human origin.

PRODUCT

Each vial contains 100 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-102464 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

CTDSPL (P-12) is recommended for detection of CTDSPL of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with family members CTDSP1 or CTDSPL2.

Suitable for use as control antibody for CTDSPL siRNA (h): sc-78502, CTDSPL shRNA Plasmid (h): sc-78502-SH and CTDSPL shRNA (h) Lentiviral Particles: sc-78502-V.

Molecular Weight of CTDSPL: 31 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.