



## DNAH1 (P-14): sc-102482

### BACKGROUND

Dyneins are multisubunit, high molecular weight ATPases that interact with microtubules to generate force by converting the chemical energy of ATP into the mechanical energy of movement. Axonemal Dynein motors contain one to three non-identical heavy chains and cause a sliding of microtubules in the axonemes of cilia and flagella in a mechanism necessary for cilia to beat and propel the cell. DNAH1 (dynein heavy chain 1, axonemal), also known as heat shock regulated protein 1 or ciliary dynein heavy chain 1, is a 4,330 amino acid protein consisting of at least two heavy chains and several intermediate and light chains. Mutations in the gene encoding DNAH1 may be a cause of primary ciliary dyskinesia, also known as Kartagener Syndrome, which is characterized by chronic recurrent respiratory infections due to defective cilia action in the respiratory tract. There are three isoforms of DNAH1 that exist as a result of alternative splicing events.

### REFERENCES

- Vaughan, K.T., Mikami, A., Paschal, B.M., Holzbaur, E.L., Hughes, S.M., Echeverri, C.J., Moore, K.J., Gilbert, D.J., Copeland, N.G., Jenkins, N.A. and Vallee, R.B. 1996. Multiple mouse chromosomal loci for dynein-based motility. *Genomics* 36: 29-38.
- Chapelin, C., Duriez, B., Magnino, F., Goossens, M., Escudier, E. and Amselem, S. 1997. Isolation of several human axonemal dynein heavy chain genes: genomic structure of the catalytic site, phylogenetic analysis and chromosomal assignment. *FEBS Lett.* 412: 325-330.
- Neesen, J., Koehler, M.R., Kirschner, R., Steinlein, C., Kreutzberger, J., Engel, W. and Schmid, M. 1997. Identification of dynein heavy chain genes expressed in human and mouse testis: chromosomal localization of an axonemal dynein gene. *Gene* 200: 193-202.
- Yagi, T. 2000. ADP-dependent microtubule translocation by flagellar inner-arm dyneins. *Cell Struct. Funct.* 25: 263-267.
- Maiti, A.K., Mattei, M.G., Jorissen, M., Volz, A., Zeigler, A. and Bouvagnet, P. 2000. Identification, tissue specific expression, and chromosomal localization of several human dynein heavy chain genes. *Eur. J. Hum. Genet.* 8: 923-932.
- Bartoloni, L., Blouin, J.L., Maiti, A.K., Sainsbury, A., Rossier, C., Gehrig, C., She, J.X., Marron, M.P., Lander, E.S., Meeks, M., Chung, E., Armengot, M., Jorissen, M., Scott, H.S., Delozier-Blanchet, C.D., Gardiner, R.M. and Antonarakis, S.E. 2001. Axonemal  $\beta$  heavy chain dynein DNAH9: cDNA sequence, genomic structure, and investigation of its role in primary ciliary dyskinesia. *Genomics* 72: 21-33.
- Neesen, J., Kirschner, R., Ochs, M., Schmiedl, A., Habermann, B., Mueller, C., Holstein, A.F., Nuesslein, T., Adham, I. and Engel, W. 2001. Disruption of an inner arm dynein heavy chain gene results in asthenozoospermia and reduced ciliary beat frequency. *Hum. Mol. Genet.* 10: 1117-1128.
- Online Mendelian Inheritance in Man, OMIM™. 2004. Johns Hopkins University, Baltimore, MD. MIM Number: 603332. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

### CHROMOSOMAL LOCATION

Genetic locus: DNAH1 (human) mapping to 3p21.1; Dnahc1 (mouse) mapping to 14 B.

### SOURCE

DNAH1 (P-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of DNAH1 of human origin.

### PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-102482 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

### APPLICATIONS

DNAH1 (P-14) is recommended for detection of DNAH1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other DNAH family members.

Suitable for use as control antibody for DNAH1 siRNA (h): sc-78027, DNAH1 siRNA (m): sc-143075, DNAH1 shRNA Plasmid (h): sc-78027-SH, DNAH1 shRNA Plasmid (m): sc-143075-SH, DNAH1 shRNA (h) Lentiviral Particles: sc-78027-V and DNAH1 shRNA (m) Lentiviral Particles: sc-143075-V.

Molecular Weight of DNAH1: 494 kDa.

### RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

### STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.