

NOS1 (K-20): sc-1025

BACKGROUND

Nitric oxide (NO) has a broad range of biological activities and has been implicated in signaling pathways in phylogenetically diverse species. Nitric oxide synthases (NOSs), the enzymes responsible for synthesis of NO, contain an N-terminal oxygenase domain and a C-terminal reductase domain. NOS activity requires homodimerization as well as three cosubstrates (L-arginine, NADPH and O₂) and five cofactors or prosthetic groups (FAD, FMN, calmodulin, tetrahydrobiopterin and heme). Several distinct NOS isoforms have been described and been shown to represent the products of three distinct genes. These include two constitutive Ca²⁺/CaM-dependent forms of NOS, including NOS1 (also designated ncNOS) whose activity was first identified in neurons, and NOS3 (also designated eNOS), first identified in endothelial cells. The inducible form of NOS, NOS2 (also designated iNOS), is Ca²⁺-independent and is expressed in a broad range of cell types.

CHROMOSOMAL LOCATION

Genetic locus: NOS1 (human) mapping to 12q24.22; Nos1 (mouse) mapping to 5 F.

SOURCE

NOS1 (K-20) is an affinity purified rabbit polyclonal antibody raised against a peptide mapping at the N-terminus of NOS1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-1025 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

NOS1 (K-20) is recommended for detection of NOS1 (ncNOS) of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:50-1:500), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:25, dilution range 1:25-1:250) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

NOS1 (K-20) is also recommended for detection of NOS1 (ncNOS) in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for NOS1 siRNA (h): sc-29416, NOS1 siRNA (m): sc-36091, NOS1 shRNA Plasmid (h): sc-29416-SH, NOS1 shRNA Plasmid (m): sc-36091-SH, NOS1 shRNA (h) Lentiviral Particles: sc-29416-V and NOS1 shRNA (m) Lentiviral Particles: sc-36091-V.

Molecular Weight of NOS1: 155 kDa.

Positive Controls: mouse brain extract: sc-2253, A-673 cell lysate: sc-2414 or rat brain extract: sc-2392.


STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.


RESEARCH USE

For research use only, not for use in diagnostic procedures.

DATA



NOS1 (K-20): sc-1025. Western blot analysis of NOS1 expression in mouse brain tissue extract.



NOS1 (K-20): sc-1025. Immunofluorescence staining of methanol-fixed A-673 cells showing cytoplasmic staining.

SELECT PRODUCT CITATIONS

- Lin, C.S., et al. 1998. Analysis of neuronal nitric oxide synthase isoform expression and identification of human nNOS-µ. *Biochem. Biophys. Res. Commun.* 253: 388-394.
- Tripathi, V. and Krishna, A. 2008. Changes in nitric oxide (NO) synthase isoforms and NO in the ovary of *Heteropneustes fossilis* (Bloch.) during the reproductive cycle. *J. Endocrinol.* 199: 307-316.
- Brahma, M.K., et al. 2009. The neuronal apoptotic death in global cerebral ischemia in gerbil: important role for sodium channel modulator. *J. Neurosci. Res.* 87: 1400-1411.
- Uytenbroek, L., et al. 2010. Neurochemical coding of enteric neurons in adult and embryonic zebrafish (*Danio rerio*). *J. Comp. Neurol.* 518: 4419-4438.
- Sánchez-Blázquez, P., et al. 2010. Mu-opioid receptors transiently activate the Akt-nNOS pathway to produce sustained potentiation of PKC-mediated NMDAR-CaMKII signaling. *PLoS ONE* 5: e11278.
- Saluja, R., et al. 2010. Ultrastructural immunogold localization of nitric oxide synthase isoforms in rat and human eosinophils. *Cell Tissue Res.* 340: 381-388.
- Mizunoya, W., et al. 2011. Nitric-oxide donors improve prednisone effects on muscular dystrophy in the mdx mouse diaphragm. *Am. J. Physiol., Cell Physiol.* 300: C1065-C1077.
- Vieira, R.P., et al. 2011. Airway epithelium mediates the anti-inflammatory effects of exercise on asthma. *Respir. Physiol. Neurobiol.* 175: 383-389.
- Gruol, D.L., et al. 2011. Neuroadaptive changes in cerebellar neurons induced by chronic exposure to IL-6. *J. Neuroimmunol.* 239: 28-36.


 MONOS
Satisfaction
Guaranteed

Try **NOS1 (A-11): sc-5302** or **NOS1 (H-7): sc-55521**, our highly recommended monoclonal alternatives to NOS1 (K-20). Also, for AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647 conjugates, see **NOS1 (A-11): sc-5302**.