

GHRH (V-17): sc-10280

BACKGROUND

Growth hormone-releasing hormone (GHRH, also designated somatoliberin or GRF) is a member of the GRF superfamily of structurally related peptide hormones. Members of the GRF superfamily have amphiphilic α -helical secondary structures as their preferred bioactive conformations. GHRH exerts its action through high-affinity GHRH receptors (GHRH-R) present in the anterior pituitary. GHRH also functions as an autocrine/paracrine growth factor for small cell lung carcinoma. GHRH stimulates secretion and synthesis of growth hormone, causes somatotroph proliferation, and may have direct actions in fetal/placental development, reproduction and immune function. The GRF superfamily includes vasoactive intestinal peptide, pituitary adenylate cyclase-activating polypeptide, secretin and glucagon. The gene encoding GHRH maps to human chromosome 20q11.2.

REFERENCES

1. Campbell, R.M., et al. 1992. Evolution of the growth hormone-releasing factor (GRF) family of peptides. *Growth Regul.* 2: 175-191.
2. Mayo, K.E. 1992. Molecular cloning and expression of a pituitary-specific receptor for growth hormone-releasing hormone. *Mol. Endocrinol.* 6: 1734-1744.
3. Gaylinn, B.D., et al. 1993. Molecular cloning and expression of a human anterior pituitary receptor for growth hormone-releasing hormone. *Mol. Endocrinol.* 7: 77-84.
4. Perez Jurado, L.A., et al. 1994. Genetic mapping of the human growth hormone-releasing factor gene (GHRF) using two intragenic polymorphisms detected by PCR amplification. *Genomics* 20: 132-134.
5. Kiaris, H., et al. 1999. Growth hormone-releasing hormone: an autocrine growth factor for small cell lung carcinoma. *Proc. Natl. Acad. Sci. USA* 96: 14894-14898.

CHROMOSOMAL LOCATION

Genetic locus: GHRH (human) mapping to 20q11.2.

SOURCE

GHRH (V-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of GHRH of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-10280 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

GHRH (V-17) is recommended for detection of precursor and mature GHRH and VIP of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for GHRH siRNA (h): sc-39519, GHRH shRNA Plasmid (h): sc-39519-SH and GHRH shRNA (h) Lentiviral Particles: sc-39519-V.

Molecular Weight of GHRH: 12 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

SELECT PRODUCT CITATIONS

1. Havt, A., et al. 2005. The expression of the pituitary growth hormone-releasing hormone receptor and its splice variants in normal and neoplastic human tissues. *Proc. Natl. Acad. Sci. USA* 102: 17424-17429.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.