



# Peregrin (S-14): sc-103111

## BACKGROUND

Peregrin, also called BRPF1 or BR140, is a 1,214 amino acid bromodomain and PHD finger-containing protein. Localized to the nucleus and expressed ubiquitously with the highest levels of expression in testis and spermatogonia, Peregrin is a zinc-finger protein that is thought to be a transcriptional activator. Peregrin contains one C-terminal PWWP domain and, because of its similarity with transcriptional coactivators such as TAF II p250, is thought to be a DNA-binding protein. The gene encoding Peregrin is disrupted in myeloid leukemia, suggesting a possible role in carcinogenesis. Due to alternative splicing, two isoforms of the Peregrin protein exist.

## REFERENCES

1. Thompson, K.A., Wang, B., Argraves, W.S., Giancotti, F.G., Schranck, D.P. and Ruoslahti, E. 1994. BR140, a novel zinc-finger protein with homology to the TAF250 subunit of TFIID. *Biochem. Biophys. Res. Commun.* 198: 1143-1152.
2. Chaplin, T., Ayton, P., Bernard, O.A., Saha, V., Della Valle, V., Hillion, J., Gregorini, A., Lillington, D., Berger, R. and Young, B.D. 1995. A novel class of zinc-finger/leucine zipper genes identified from the molecular cloning of the t(10;11) translocation in acute leukemia. *Blood* 85: 1435-1441.
3. Gregorini, A., Sahin, F.I., Lillington, D.M., Meerabux, J., Saha, V., McCullagh, P., Bocci, M., Menevse, S., Papa, S. and Young, B.D. 1997. Gene BR140, which is related to AF10 and AF17, maps to chromosome band 3p25. *Genes Chromosomes Cancer*. 17: 269-272.
4. McCullagh, P., Chaplin, T., Meerabux, J., Grenzeliass, D., Lillington, D., Poulson, R., Gregorini, A., Saha, V. and Young, B.D. 2000. The cloning, mapping and expression of a novel gene, BRL, related to the AF10 leukaemia gene. *Oncogene* 18: 7442-7452.
5. Chamberlin, H.M. and Thomas, J.H. 2000. The bromodomain protein LIN-49 and trithorax-related protein LIN-59 affect development and gene expression in *Caenorhabditis elegans*. *Development* 127: 713-723.
6. Stec, I., Nagl, S.B., van Ommen, G.J. and den Dunnen, J.T. 2000. The PWWP domain: a potential protein-protein interaction domain in nuclear proteins influencing differentiation? *FEBS Lett.* 473: 1-5.
7. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 602410. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

## CHROMOSOMAL LOCATION

Genetic locus: BRPF1 (human) mapping to 3p25.3; Brpf1 (mouse) mapping to 6 E3.

## SOURCE

Peregrin (S-14) is an affinity purified rabbit polyclonal antibody raised against a peptide mapping near the N-terminus of Peregrin of human origin.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PRODUCT

Each vial contains 100 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-103111 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-103111 X, 100 µg/0.1 ml.

## APPLICATIONS

Peregrin (S-14) is recommended for detection of Peregrin of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Peregrin siRNA (h): sc-78127, Peregrin shRNA Plasmid (h): sc-78127-SH and Peregrin shRNA (h) Lentiviral Particles: sc-78127-V.

Peregrin (N-16) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Molecular Weight of Peregrin: 150 kDa.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.