

scotin (P-12): sc-103189

BACKGROUND

Scotin (protein shisha-5), also known as putative NF κ B-activating protein 120, is a 240 amino acid single-pass type I membrane protein that localizes to the endoplasmic reticulum and nucleus. Scotin belongs to the shisha protein family and contains a proline-rich domain. Both caspase-dependent and p53/TP53-dependent apoptosis appear to be induced by scotin. Scotin is abundant in murine spleen and thymus tissue. The gene encoding scotin maps to human chromosome 3, which houses over 1,100 genes, including a chemokine receptor (CKR) gene cluster and a variety of human cancer-related gene loci. Key tumor suppressing genes on chromosome 3 include those that encode the apoptosis mediator RASSF1, the cell migration regulator HYAL1 and the angiogenesis suppressor SEMA3B. Marfan syndrome, porphyria, von Hippel-Lindau syndrome, osteogenesis imperfecta and Charcot-Marie-Tooth disease are a few of the numerous genetic diseases associated with chromosome 3.

REFERENCES

1. Bourdon, J.C., Renzing, J., Robertson, P.L., Fernandes, K.N. and Lane, D.P. 2002. Scotin, a novel p53-inducible proapoptotic protein located in the ER and the nuclear membrane. *J. Cell Biol.* 158: 235-246.
2. Muzny, D.M., Scherer, S.E., Kaul, R., Wang, J., Yu, J., Sudbrak, R., Buhay, C.J., Chen, R., Cree, A., Ding, Y., Dugan-Rocha, S., Gill, R., Gunaratne, P., Harris, R.A., Hawes, A.C., Hernandez, J., Hodgson, A.V., et al. 2006. The DNA sequence, annotation and analysis of human chromosome 3. *Nature* 440: 1194-1198.
3. Draeby, I., Woods, Y.L., la Cour, J.M., Mollerup, J., Bourdon, J.C. and Berchtold, M.W. 2007. The calcium binding protein ALG-2 binds and stabilizes scotin, a p53-inducible gene product localized at the endoplasmic reticulum membrane. *Arch. Biochem. Biophys.* 467: 87-94.
4. Zocchi, L., Bourdon, J.C., Codispoti, A., Knight, R., Lane, D.P., Melino, G. and Terrinoni, A. 2008. Scotin: A new p63 target gene expressed during epidermal differentiation. *Biochem. Biophys. Res. Commun.* 367: 271-276.
5. Gupta, R.K., Tripathi, R., Naidu, B.J., Srinivas, U.K. and Shashidhara, L.S. 2008. Cell cycle regulation by the pro-apoptotic gene scotin. *Cell Cycle* 7: 2401-2408.
6. Ghosh, S., Ghosh, A., Maiti, G.P., Alam, N., Roy, A., Roy, B., Roychoudhury, S. and Panda, C.K. 2008. Alterations of 3p21.31 tumor suppressor genes in head and neck squamous cell carcinoma: Correlation with progression and prognosis. *Int. J. Cancer.* 123: 2594-2604.

CHROMOSOMAL LOCATION

Genetic locus: SHISA5 (human) mapping to 3p21.31, LOC646056 (human) mapping to Xq13.2.

SOURCE

scotin (P-12) is an affinity purified rabbit polyclonal antibody raised against a peptide mapping within a cytoplasmic domain of scotin of human origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-103189 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

scotin (P-12) is recommended for detection of scotin and LOC646056 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
 Satisfation
 Guaranteed

Try **scotin (C-7): sc-390725**, our highly recommended monoclonal alternative to scotin (P-12).