

CCL27 (H-16): sc-10323

BACKGROUND

Cutaneous T cell-attracting chemokine, CCL27 (also designated CTACK, ILC and ALP) is a member of the chemokine superfamily and the subfamily of β or C-C chemokines that binds chemokine receptor CCR10 (GPR-2). Chemokines are a superfamily of small secreted proteins that attract their targets by interacting with G protein-coupled receptors expressed on the migrating cell. CCL27 and CCR10 are involved in skin homeostasis and inflammatory response. CCL27 is found in human keratinocytes and mouse epidermis; CCR10 is expressed by melanocytes, dermal fibroblasts and dermal microvascular endothelial cells and in T cells, as well as in skin-derived Langerhans cells. CCL27 is involved in the preferential migration and recruitment of cutaneous lymphocyte-associated antigen CLA⁺ memory lymphocytes.

REFERENCES

1. Baggiolini, M. 1998. Chemokines and leukocyte traffic. *Nature* 392: 565-568.
2. Morales, J., et al. 1999. CTACK, a skin-associated chemokine that preferentially attracts skin-homing memory T cells. *Proc. Natl. Acad. Sci. USA* 96: 14470-14475.
3. Ishikawa-Mochizuki, I., et al. 1999. Molecular cloning of a novel C-C chemokine, interleukin-11 receptor α -locus chemokine (ILC), which is located on chromosome 9p13 and a potential homologue of a C-C chemokine encoded by mollusum contagiosum virus. *FEBS Lett.* 460: 544-548.
4. Zaballos, A., et al. 1999. CCL27, the human homologue of murine ALP chemokine. Submitted to EMBL/GenBank/DBJ database.
5. Hromas, R., et al. 1999. Isolation of ALP, a novel divergent murine C-C chemokine with a unique carboxy-terminal extension. *Biochem. Biophys. Res. Commun.* 258: 737-740.
6. Zlotnik, A., et al. 1999. Recent advances in chemokines and chemokine receptors. *Crit. Rev. Immunol.* 19: 1-47.
7. Homey, B., et al. 2000. Cutting edge: the orphan chemokine receptor G protein-coupled receptor-2 (GPR-2, CCR10) binds the skin-associated chemokine CCL27. *J. Immunol.* 164: 3465-3470.

CHROMOSOMAL LOCATION

Genetic locus: CCL27 (human) mapping to 9p13.

SOURCE

CCL27 (H-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of CCL27 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-10323 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

CCL27 (H-16) is recommended for detection of CCL27 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for CCL27 siRNA (h): sc-39346, CCL27 shRNA Plasmid (h): sc-39346-SH and CCL27 shRNA (h) Lentiviral Particles: sc-39346-V.

Molecular Weight of CCL27: 10 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **CCL27 (G-6): sc-390112**, our highly recommended monoclonal alternative to CCL27 (H-16).