

UMPS (S-16): sc-103313

BACKGROUND

Uridine 5'-monophosphate synthase (UMPS) catalyzes the last two steps of the pyrimidine biosynthetic pathway. Unlike prokaryotes, UMPS in eukaryotes combines the orotate phosphoribosyltransferase and the orotidine-5'-monophosphate (OMP) decarboxylase activities into a single protein. The union of these two enzymes is thought to stabilize the catalytic centers due to the low molar concentration of the protein in mammalian cells. Loss of either enzymatic activity results in hereditary orotic aciduria, a rare autosomal recessive disorder characterized by retarded growth, anemia, and excessive urinary excretion of orotic acid. Two isoforms of UMPS exist as a result of alternative splicing events.

CHROMOSOMAL LOCATION

Genetic locus: UMPS (human) mapping to 3q21.2; Umps (mouse) mapping to 16 B3.

SOURCE

UMPS (S-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of UMPS of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-103313 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures

APPLICATIONS

UMPS (S-16) is recommended for detection of UMPS of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

UMPS (S-16) is also recommended for detection of UMPS in additional species, including equine, canine, bovine and avian.

Suitable for use as control antibody for UMPS siRNA (h): sc-78096, UMPS siRNA (m): sc-154917, UMPS shRNA Plasmid (h): sc-78096-SH, UMPS shRNA Plasmid (m): sc-154917-SH, UMPS shRNA (h) Lentiviral Particles: sc-78096-V and UMPS shRNA (m) Lentiviral Particles: sc-154917-V.


Molecular Weight of UMPS: 52/33 kDa.

Positive Controls: UMPS (h2): 293T Lysate: sc-178109, MOLT-4 cell lysate: sc-2233 or IMR-32 cell lysate: sc-2409.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



UMPS (S-16): sc-103313. Western blot analysis of UMPS expression in non-transfected 293T: sc-117752 (A), human UMPS transfected 293T: sc-178109 (B), MOLT-4 (C) and IMR-32 (D) whole cell lysates.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **UMPS (A-9): sc-398086** or **UMPS (3H6): sc-135596**, our highly recommended monoclonal alternatives to UMPS (S-16).