



Arylsulfatase H (S-16): sc-103397

BACKGROUND

Sulfatases hydrolyze sulfate esters from sulfated steroids, carbohydrates, proteoglycans and glycolipids. They are involved in hormone biosynthesis, modulation of cell signaling and degradation of macromolecules. Arylsulfatase H, also known as ARSH, is a 562 amino acid protein that belongs to the sulfatase family of bone and cartilage matrix proteins. Localized to the plasma membrane, Arylsulfatase H uses calcium as a cofactor to hydrolyze sulfate esters. The gene encoding Arylsulfatase D maps to human chromosome X, which contains nearly 153 million base pairs and houses over 1,000 genes. In conjunction with chromosome Y, chromosome X is responsible for sex determination. There are a number of conditions related to an abnormal number and combination of sex chromosomes, some of which include Turner's syndrome, color blindness, hemophilia and Duchenne muscular dystrophy.

REFERENCES

1. Chant, J. and Stowers, L. 1995. GTPase cascades choreographing cellular behavior: movement, morphogenesis, and more. *Cell* 81: 1-4.
2. Franco, B., Meroni, G., Parenti, G., Levilliers, J., Bernard, L., Gebbia, M., Cox, L., Maroteaux, P., Sheffield, L. and Rappold, G.A. 1995. A cluster of sulfatase genes on Xp22.3: mutations in chondrodysplasia punctata (CDPX) and implications for warfarin embryopathy. *Cell* 81: 15-25.
3. Meroni, G., Franco, B., Archidiacono, N., Messali, S., Andolfi, G., Rocchi, M. and Ballabio, A. 1996. Characterization of a cluster of sulfatase genes on Xp22.3 suggests gene duplications in an ancestral pseudoautosomal region. *Hum. Mol. Genet.* 5: 423-431.
4. Puca, A.A., Zollo, M., Repetto, M., Andolfi, G., Guffanti, A., Simon, G., Ballabio, A. and Franco, B. 1997. Identification by shotgun sequencing, genomic organization, and functional analysis of a fourth arylsulfatase gene (ARSF) from the Xp22.3 region. *Genomics* 42: 192-199.
5. Dooley, T.P., Haldeman-Cahill, R., Joiner, J. and Wilborn, T.W. 2000. Expression profiling of human sulfotransferase and sulfatase gene superfamilies in epithelial tissues and cultured cells. *Biochem. Biophys. Res. Commun.* 277: 236-245.
6. Urbitsch, P., Salzer, M.J., Hirschmann, P. and Vogt, P.H. 2000. Arylsulfatase D gene in Xp22.3 encodes two protein isoforms. *DNA Cell Biol.* 19: 765-773.
7. Sardiello, M., Annunziata, I., Roma, G. and Ballabio, A. 2005. Sulfatases and sulfatase modifying factors: an exclusive and promiscuous relationship. *Hum. Mol. Genet.* 14: 3203-3217.
8. Online Mendelian Inheritance in Man, OMIM™. 2006. Johns Hopkins University, Baltimore, MD. MIM Number: 300586. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
9. Lin, Y.F., Yang, J. and Rosen, B.P. 2007. ArsD residues Cys12, Cys13, and Cys18 form an As(III)-binding site required for arsenic metallochaperone activity. *J. Biol. Chem.* 282: 16783-16791.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

CHROMOSOMAL LOCATION

Genetic locus: ARSH (human) mapping to Xp22.33.

SOURCE

Arylsulfatase H (S-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of Arylsulfatase H of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-103397 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Arylsulfatase H (S-16) is recommended for detection of Arylsulfatase H of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other AS family members.

Suitable for use as control antibody for Arylsulfatase H siRNA (h): sc-105094, Arylsulfatase H shRNA Plasmid (h): sc-105094-SH and Arylsulfatase H shRNA (h) Lentiviral Particles: sc-105094-V.

Molecular Weight of Arylsulfatase H: 63 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.