

## cGKII (N-16): sc-10345

### BACKGROUND

cGKII (cGMP-dependent protein kinase type II) is a major receptor of intracellular cGMP and mediates a plethora of physiological responses. cGKII contains a conserved leucine zipper motif at the amino-terminus. It is expressed in small intestine, colon, prostate, and human brain tissues, and the cGKII gene maps to chromosome 4q21.1. cGKII has been shown to regulate the ion transport system in the intestine. Myristoylation of the penultimate glycine in cGKII appears to be essential for directing cGKII to the membrane, since cGKII is devoid of any hydrophobic transmembrane domains. The translocation of cGKII from the cytosol to the membrane allows it to function properly in regulating intestinal ion transport.

### REFERENCES

1. Uhler, M.D. 1993. Cloning and expression of a novel cyclic GMP-dependent protein kinase from mouse brain. *J. Biol. Chem.* 268: 13586-13591.
2. Gamm, D.M., et al. 1995. The type II isoform of cGMP-dependent protein kinase is dimeric and possesses regulatory and catalytic properties distinct from the type I isoforms. *J. Biol. Chem.* 270: 27380-27388.
3. Vaandrager, A.B., et al. 1996. Signalling by cGMP-dependent protein kinases. *Mol. Cell. Biochem.* 157: 23-30.
4. Orstavik, S., et al. 1996. Molecular cloning, cDNA structure, and chromosomal localization of the human type II cGMP-dependent protein kinase. *Biochem. Biophys. Res. Commun.* 220: 759-765.
5. Vaandrager, A.B., et al. 1996. N-terminal myristoylation is required for membrane localization of cGMP-dependent protein kinase type II. *J. Biol. Chem.* 271: 7025-7029.
6. Wagner, C., et al. 1998. Role of cGMP-kinase II in the control of renin secretion and renin expression. *J. Clin. Invest.* 102: 1576-1582.

### CHROMOSOMAL LOCATION

Genetic locus: PRKG2 (human) mapping to 4q21.21.

### SOURCE

cGKII (N-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of cGKII of human origin.

### PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-10345 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

### STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### APPLICATIONS

cGKII (N-16) is recommended for detection of cGKII of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

cGKII (N-16) is also recommended for detection of cGKII in additional species, including equine.


Suitable for use as control antibody for cGKII siRNA (h): sc-38974, cGKII shRNA Plasmid (h): sc-38974-SH and cGKII shRNA (h) Lentiviral Particles: sc-38974-V.

Molecular Weight of cGKII: 86 kDa.

### RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 3) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

### DATA



cGKII (N-16): sc-10345. Immunoperoxidase staining of formalin fixed, paraffin-embedded human small intestine tissue showing membrane and cytoplasmic staining of glandular cells.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.

**MONOS**  
Satisfaction  
Guaranteed

Try **cGKII (D-3): sc-393126** or **cGKII (E-7): sc-390926**, our highly recommended monoclonal alternatives to cGKII (N-16).