



Dnmt3L (S-14): sc-103480

BACKGROUND

Methylation at the 5'-position of cytosine is the only known naturally occurring covalent modification of the mammalian genome. DNA methylation requires the enzymatic activity of DNA 5-cytosine methyltransferase (Dnmt) proteins, which catalyze the transfer of a methyl group from S-adenosyl methionine to the 5'-position of cytosines, thereby repressing expression of the target gene. Dnmt3L (DNA (cytosine-5)-methyltransferase 3-like) is a 387 amino acid protein that contains one ADD-type zinc finger and is a member of the Dnmt family. Localized to the nucleus and expressed at low levels in thymus, testis and ovary, Dnmt3L does not exhibit DNA methyltransferase activity, but is able to stimulate *de novo* methylation by Dnmt3 and is thought to play a key role in the establishment of genomic imprints. Additionally, Dnmt3L interacts with histone deacetylase 1 (HDAC1) and, through this interaction, mediates transcriptional repression. Multiple isoforms of Dnmt3L exist due to alternative splicing events.

REFERENCES

1. Yoder, J.A., et al. 1997. DNA (cytosine-5)-methyltransferases in mouse cells and tissues. Studies with a mechanism-based probe. *J. Mol. Biol.* 270: 385-395.
2. Walsh, C.P. and Bestor, T.H. 1999. Cytosine methylation and mammalian development. *Genes Dev.* 13: 26-34.
3. Hsieh, C.L. 1999. *In vivo* activity of murine *de novo* methyltransferases, Dnmt3a and Dnmt3b. *Mol. Cell. Biol.* 19: 8211-8218.
4. Fuks, F., et al. 2000. DNA methyltransferase Dnmt1 associates with histone deacetylase activity. *Nat. Genet.* 24: 88-91.
5. Aapola, U., et al. 2000. Isolation and initial characterization of a novel zinc finger gene, Dnmt3L, on 21q22.3, related to the cytosine-5-methyltransferase 3 gene family. *Genomics* 65: 293-298.
6. Suetake, I., et al. 2006. Stimulation effect of Dnmt3L on the DNA methylation activity of Dnmt3a2. *J. Biochem.* 140: 553-559.
7. Ooi, S.K., et al. 2007. Dnmt3L connects unmethylated Lysine 4 of Histone H3 to *de novo* methylation of DNA. *Nature* 448: 714-717.
8. Jia, D., et al. 2007. Structure of Dnmt3a bound to Dnmt3L suggests a model for *de novo* DNA methylation. *Nature* 449: 248-251.
9. Gokul, G., et al. 2007. DNA methylation profile at the Dnmt3L promoter: a potential biomarker for cervical cancer. *Epigenetics* 2: 80-85.

CHROMOSOMAL LOCATION

Genetic locus: Dnmt3l (mouse) mapping to 10 C1.

SOURCE

Dnmt3L (S-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Dnmt3L of mouse origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-103480 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Dnmt3L (S-14) is recommended for detection of Dnmt3L of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other Dnmt family members.

Suitable for use as control antibody for Dnmt3L siRNA (m): sc-37762, Dnmt3L shRNA Plasmid (m): sc-37762-SH and Dnmt3L shRNA (m) Lentiviral Particles: sc-37762-V.

Molecular Weight of Dnmt3L: 43 kDa.

Positive Controls: mouse testis extract: sc-2405.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.