

# IGSF4B (Y-15): sc-103560

## BACKGROUND

Immunoglobulin superfamily member 4B (cell adhesion molecule 3, Nectin-like protein 1) is a nectin family gene product that contains two Ig-like C2-type (immunoglobulin-like) domains and an Ig-like V-type (immunoglobulin-like) domain. IGSF4B is a single-pass type I membrane protein that localizes to cell-cell interacting sites along the plasma membrane. IGSF4B functions as a cell adhesion molecule at cell-cell junctions. IGSF4B can be found as a homodimer and has a dual calcium-independent adhesion function; a homophilic cell-cell interaction and a heterophilic cell-cell interaction involving IGSF4, Nectin 1 and Nectin 3. IGSF4B may also interact with EPB41L1 which has a potential for regulating structure or function of cell-cell junctions. By similarity, Isoform 1 is expressed in adult and fetal brain. Isoform 2 is highly expressed in the cerebellum but is also weakly expressed in placenta. IGSF4B has been found to be markedly over-expressed in glioma cell lines and prostate cancer cell lines.

## REFERENCES

1. Fukuhara, H., et al. 2001. Isolation of the TSL1 and TSL2 genes, members of the tumor suppressor TSLC1 gene family encoding transmembrane proteins. *Oncogene* 20: 5401-5407.
2. Dong, X., et al. 2006. Crystal structure of the V domain of human Nectin-like molecule-1/Syncam3/Tsl1/IGSF4B, a neural tissue-specific immunoglobulin-like cell-cell adhesion molecule. *J. Biol. Chem.* 281: 10610-10617.
3. Gao, J., et al. 2008. Nectin-like molecule 1 is a glycoprotein with a single N-glycosylation site at N290K which influences its adhesion activity. *Biochim. Biophys. Acta* 1778: 1429-1435.
4. Chen, T., et al. 2008. Role of cell adhesion molecules Necl1 in synaptogenesis in primary cultured rat neurons. *Zhongguo Yi Xue Ke Xue Yuan Xue Bao* 30: 275-279.
5. Gao, J., et al. 2008. Effect of NECL1 on the proliferation of T98G glioma cell line. *Zhongguo Yi Xue Ke Xue Yuan Xue Bao* 30: 280-283.
6. Gao, J., et al. 2009. Loss of NECL1, a novel tumor suppressor, can be restored in glioma by HDAC inhibitor-Trichostatin A through Sp1 binding site. *Glia* 57: 989-999.
7. Raveh, S., et al. 2009. The cell adhesion Nectin-like molecules (Necl) 1 and 4 suppress the growth and tumorigenic ability of colon cancer cells. *J. Cell. Biochem.* 108: 326-336.
8. Spiegel, I., et al. 2009. A novel method for isolating Schwann cells using the extracellular domain of Necl1. *J. Neurosci. Res.* 87: 3288-3296.
9. Yin, B., et al. 2009. Neural adhesion molecule NECL1 inhibits migration, invasion, and potentially induces differentiation of glioma cell. *Zhongguo Yi Xue Ke Xue Yuan Xue Bao* 31: 669-673.

## CHROMOSOMAL LOCATION

Genetic locus: CADM3 (human) mapping to 1q23.2; Cadm3 (mouse) mapping to 1 H3.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## SOURCE

IGSF4B (Y-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an extracellular domain of IGSF4B of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-103560 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

IGSF4B (Y-15) is recommended for detection of IGSF4B of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with family members IGSF4C or IGSF4D.


IGSF4B (Y-15) is also recommended for detection of IGSF4B in additional species, including canine, bovine and porcine.

Suitable for use as control antibody for IGSF4B siRNA (h): sc-88048, IGSF4B siRNA (m): sc-105563, IGSF4B shRNA Plasmid (h): sc-88048-SH, IGSF4B shRNA Plasmid (m): sc-105563-SH, IGSF4B shRNA (h) Lentiviral Particles: sc-88048-V and IGSF4B shRNA (m) Lentiviral Particles: sc-105563-V.

Molecular Weight of IGSF4B: 44 kDa.

Positive Controls: SK-MEL-28 cell lysate: sc-2236 or KNRK whole cell lysate: sc-2214.

## DATA



IGSF4B (Y-15): sc-103560. Western blot analysis of IGSF4B expression in SK-MEL-28 (A) and KNRK (B) whole cell lysates.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.