

L-type Ca⁺⁺ CP α 1C/D/F/S (G-14): sc-103590

BACKGROUND

L-type (long lasting current) voltage-dependent calcium channels are composed of four subunits, designated α 1, β , γ and α 2/ δ , all of which work together to mediate neurotransmitter release. The α subunit contains a tetrameric association of four domains each containing a series of six transmembrane α -helical segments, numbered S1-S6, which are connected by both intracellular and extracellular loops. The α subunit is comprised of an ion-conducting pore, which determines the main characteristics of the cation channel complex, including ion selectivity, voltage sensitivity and pharmacology, and binding characteristics for endogenous and exogenous ligands. There are ten genes in the human genome that encode pore-forming α 1 subunits of voltage-gated calcium channels. In combination with accessory subunits, these ten α 1 subunits, which include L-type Ca⁺⁺ CP α 1C, D, F and S, must mediate diverse functions such as intracellular calcium homeostasis, regulation of gene expression and coupling of membrane potential changes to various downstream processes like neurotransmitter release or muscle contraction.

REFERENCES

1. Safayhi, H., et al. 1997. L-type calcium channels in Insulin-secreting cells: biochemical characterization and phosphorylation in RINm5F cells. *Mol. Endocrinol.* 11: 619-629.
2. Kelly, K.M., et al. 2003. Alterations in hippocampal voltage-gated calcium channel α 1 subunit expression patterns after kainate-induced status epilepticus in aging rats. *Epilepsy Res.* 57: 15-32.
3. Jurkat-Rott, K. and Lehmann-Horn, F. 2004. The impact of splice isoforms on voltage-gated calcium channel α 1 subunits. *J. Physiol.* 554: 609-619.
4. Hayashida, S., et al. 2005. Increased expression of L-type high voltage-gated calcium channel α 1 and α 2/ δ subunits in mouse brain after chronic nicotine administration. *Brain Res. Mol. Brain Res.* 135: 280-284.
5. Felix, R. 2005. Molecular regulation of voltage-gated Ca²⁺ channels. *J. Recept. Signal Transduct. Res.* 25: 57-71.
6. Fiordeliso, T., et al. 2007. Immunoreactivity to neurofilaments in the rodent anterior pituitary is associated with the expression of α 1A protein subunits of voltage-gated Ca²⁺ channels. *J. Neuroendocrinol.* 19: 870-881.
7. Dolphin, A.C. 2009. Calcium channel diversity: multiple roles of calcium channel subunits. *Curr. Opin. Neurobiol.* 19: 237-244.
8. Obermair, G.J., et al. 2010. Reciprocal interactions regulate targeting of calcium channel β subunits and membrane expression of α 1 subunits in cultured hippocampal neurons. *J. Biol. Chem.* 285: 5776-5791.
9. Zamponi, G.W., et al. 2010. Role of voltage-gated calcium channels in epilepsy. *Pflugers Arch.* 460: 395-403.

SOURCE

L-type Ca⁺⁺ CP α 1C/D/F/S (G-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within a cytoplasmic domain of L-type Ca⁺⁺ CP α 1D of human origin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-103590 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

L-type Ca⁺⁺ CP α 1C/D/F/S (G-14) is recommended for detection of L-type Ca⁺⁺ CP α 1S/1C/1D/1F of mouse and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

L-type Ca⁺⁺ CP α 1C/D/F/S (G-14) is also recommended for detection of L-type Ca⁺⁺ CP α 1S/1C/1D/1F in additional species, including equine, canine, bovine, porcine and avian.

Molecular Weight of L-type Ca⁺⁺ CP α 1C/D/F/S short form: 199 kDa.

Molecular Weight of L-type Ca⁺⁺ CP α 1C long form: 240 kDa.


Molecular Weight of L-type Ca⁺⁺ CP α 1D: 245 kDa.

Molecular Weight of L-type Ca⁺⁺ CP α 1F: 221 kDa.


Molecular Weight of L-type Ca⁺⁺ CP α 1S: 212 kDa.

Positive Controls: IMR-32 cell lysate: sc-2409.

DATA



L-type Ca⁺⁺ CP α 1C/D/F/S (G-14): sc-103590. Western blot analysis of human recombinant L-type Ca⁺⁺ CP α 1C/D/F/S fusion protein.



L-type Ca⁺⁺ CP α 1C/D/F/S (G-14): sc-103590. Western blot analysis of L-type Ca⁺⁺ CP α 1C/D/F/S expression in IMR-32 whole cell lysate.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.


 MONOS
Satisfaction
Guaranteed

Try **L-type Ca⁺⁺ CP α 1C (D-6): sc-398433** or **L-type Ca⁺⁺ CP α 1D (G-9): sc-515643**, our highly recommended monoclonal alternatives to L-type Ca⁺⁺ CP α 1C/D/F/S (G-14).