# PCDHA9 (S-13): sc-103804



The Power to Question

## **BACKGROUND**

Protocadherins are a large family of cadherin-like cell adhesion proteins that are involved in the establishment and maintenance of neuronal connections in the brain. There are three protocadherin gene clusters, designated  $\alpha,\,\beta$  and  $\gamma,$  all of which contain multiple tandemly arranged genes. PCDHA9 (protocadherin  $\alpha$  9), also known as KIAA0345, is a 950 amino acid single-pass type I membrane protein that contains six cadherin domains and is encoded by a gene which is located within the protocadherin  $\alpha$  gene cluster on human chromosome 5. Existing as multiple alternatively spliced isoforms, PCDHA9 functions as a potential calcium-dependent cell adhesion protein that may be involved in the establishment and maintenance of neuronal connections within the brain.

## **REFERENCES**

- Suzuki, S.T. 2000. Recent progress in protocadherin research. Exp. Cell Res. 261: 13-18.
- Yagi, T. and Takeichi, M. 2000. Cadherin superfamily genes: functions, genomic organization, and neurologic diversity. Genes Dev. 14: 1169-1180.
- Nollet, F., Kools, P. and van Roy, F. 2000. Phylogenetic analysis of the cadherin superfamily allows identification of six major subfamilies besides several solitary members. J. Mol. Biol. 299: 551-572.
- Wu, Q. and Maniatis, T. 2000. Large exons encoding multiple ectodomains are a characteristic feature of protocadherin genes. Proc. Natl. Acad. Sci. USA 97: 3124-3129.
- Wu, Q., Zhang, T., Cheng, J.F., Kim, Y., Grimwood, J., Schmutz, J., Dickson, M., Noonan, J.P., Zhang, M.Q., Myers, R.M. and Maniatis, T. 2001. Comparative DNA sequence analysis of mouse and human protocadherin gene clusters. Genome Res. 11: 389-404.
- 6. Noonan, J.P., Li, J., Nguyen, L., Caoile, C., Dickson, M., Grimwood, J., Schmutz, J., Feldman, M.W. and Myers, R.M. 2003. Extensive linkage disequilibrium, a common 16.7 kilobase deletion, and evidence of balancing selection in the human protocadherin  $\alpha$  cluster. Am. J. Hum. Genet. 72: 621-635.
- 7. Ribich, S., Tasic, B. and Maniatis, T. 2006. Identification of long-range regulatory elements in the protocadherin  $\alpha$  gene cluster. Proc. Natl. Acad. Sci. USA 103: 19719-19724.
- 8. Online Mendelian Inheritance in Man, OMIM™. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 606314. World Wide Web URL: http://www.ncbi.nlm.nih.gov/omim/

## CHROMOSOMAL LOCATION

Genetic locus: Pcdha9 (rat) mapping to 18p11.

### **STORAGE**

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

# **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

#### **SOURCE**

PCDHA9 (S-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of PCDHA9 of rat origin.

## **PRODUCT**

Each vial contains 200  $\mu g$  lgG in 1.0 ml of PBS with <0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-103804 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## **APPLICATIONS**

PCDHA9 (S-13) is recommended for detection of PCDHA9 of rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other PCDHA family members.

Molecular Weight of PCDHA9: 102 kDa

## **RECOMMENDED SECONDARY REAGENTS**

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

# **PROTOCOLS**

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

**Santa Cruz Biotechnology, Inc.** 1.800.457.3801 831.457.3800 fax 831.457.3801 **Europe** +00800 4573 8000 49 6221 4503 0 **www.scbt.com**