

GPR25 (N-17): sc-103969

BACKGROUND

G protein-coupled receptors (GPRs), also known as seven transmembrane receptors, heptahelical receptors or 7TM receptors, comprise a superfamily of proteins that play a role in many different stimulus-response pathways. G protein-coupled receptors translate extracellular signals into intracellular signals (G protein activation) and they respond to a variety of signaling molecules, such as hormones and neurotransmitters. GPR25 (G protein-coupled receptor 25) is a 361 amino acid multi-pass membrane protein that belongs to the G protein-coupled receptor family and functions as an orphan receptor, possibly playing a role in signal transduction throughout the cell. The gene encoding GPR25 maps to human chromosome 1, which spans 260 million base pairs, contains over 3,000 genes and comprises nearly 8% of the human genome.

REFERENCES

1. Houslay, M.D. 1992. G-protein linked receptors: a family probed by molecular cloning and mutagenesis procedures. *Clin. Endocrinol.* 36: 525-534.
2. Larhammar, D., Blomqvist, A.G. and Wahlestedt, C. 1993. The receptor revolution—multiplicity of G-protein-coupled receptors. *Drug Des. Discov.* 9: 179-188.
3. Jung, B.P., Nguyen, T., Kolakowski, L.F., Lynch, K.R., Heng, H.H., George, S.R. and O'Dowd, B.F. 1997. Discovery of a novel human G protein-coupled receptor gene (GPR25) located on chromosome 1. *Biochem. Biophys. Res. Commun.* 230: 69-72.
4. Online Mendelian Inheritance in Man, OMIM[™]. 1997. Johns Hopkins University, Baltimore, MD. MIM Number: 602174. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Ji, T.H., Grossmann, M. and Ji, I. 1998. G protein-coupled receptors. I. Diversity of receptor-ligand interactions. *J. Biol. Chem.* 273: 17299-17302.
6. Schöneberg, T., Schultz, G. and Gudermann, T. 1999. Structural basis of G protein-coupled receptor function. *Mol. Cell. Endocrinol.* 151: 181-193.
7. Lee, D.K., Nguyen, T., Lynch, K.R., Cheng, R., Vanti, W.B., Arkhitko, O., Lewis, T., Evans, J.F., George, S.R. and O'Dowd, B.F. 2001. Discovery and mapping of ten novel G protein-coupled receptor genes. *Gene* 275: 83-91.

CHROMOSOMAL LOCATION

Genetic locus: GPR25 (human) mapping to 1q32.1.

SOURCE

GPR25 (N-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an N-terminal extracellular domain of GPR25 of human origin.

PRODUCT

Each vial contains 100 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, ready P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

GPR25 (N-17) is recommended for detection of GPR25 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other GPR family members.

Suitable for use as control antibody for GPR25 siRNA (h): sc-88152, GPR25 shRNA Plasmid (h): sc-88152-SH and GPR25 shRNA (h) Lentiviral Particles: sc-88152-V.

Molecular Weight of GPR25: 39 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.