

APOBEC4 (S-16): sc-104060

BACKGROUND

APOBEC (apolipoprotein B mRNA editing enzyme, catalytic) proteins inhibit retroviruses by deaminating cytosine residues of viral RNA and DNA. APOBEC4 (apolipoprotein B mRNA editing enzyme, catalytic polypeptide-like 4), also known as C1orf169, is a 367 amino acid protein that belongs to the AID/APOBEC family of polynucleotide deoxycytidine deaminases whose primary function is to catalyze the conversion of cytidine to uridine. Expressed predominantly in testis, APOBEC4 uses zinc as a cofactor to convert cytidine to uridine, an event that is important for RNA editing. The gene encoding APOBEC4 maps to human chromosome 1, which spans 260 million base pairs, contains over 3,000 genes and comprises nearly 8% of the human genome. Chromosome 1 houses a large number of disease-associated genes, including those that are involved in familial adenomatous polyposis, Stickler syndrome, Parkinson's disease, Gaucher disease, schizophrenia and Usher syndrome. Aberrations in chromosome 1 are found in a variety of cancers, including head and neck cancer, malignant melanoma and multiple myeloma.

REFERENCES

- Rogozin, I.B., et al. 2005. APOBEC4, a new member of the AID/APOBEC family of polynucleotide (deoxy)cytidine deaminases predicted by computational analysis. *Cell Cycle* 4: 1281-1285.
- Weise, A., et al. 2005. New insights into the evolution of chromosome 1. *Cytogenet. Genome Res.* 108: 217-222.
- Coticello, S.G., et al. 2005. Evolution of the AID/APOBEC family of polynucleotide (deoxy)cytidine deaminases. *Mol. Biol. Evol.* 22: 367-377.
- Marzin, Y., et al. 2006. Chromosome 1 abnormalities in multiple myeloma. *Anticancer Res.* 26: 953-959.
- Navaratnam, N. and Sarwar, R. 2006. An overview of cytidine deaminases. *Int. J. Hematol.* 83: 195-200.
- Online Mendelian Inheritance in Man, OMIM™. 2006. Johns Hopkins University, Baltimore, MD. MIM Number: 609908. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Coticello, S.G., et al. 2007. DNA deamination in immunity: AID in the context of its APOBEC relatives. *Adv. Immunol.* 94: 37-73.

CHROMOSOMAL LOCATION

Genetic locus: Apobec4 (mouse) mapping to 1 G3.

SOURCE

APOBEC4 (S-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of APOBEC4 of mouse origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-104060 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

APOBEC4 (S-16) is recommended for detection of APOBEC4 of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other APOBEC family members.

Suitable for use as control antibody for APOBEC4 siRNA (m): sc-105082, APOBEC4 shRNA Plasmid (m): sc-105082-SH and APOBEC4 shRNA (m) Lentiviral Particles: sc-105082-V.


Molecular Weight of APOBEC4: 42 kDa.

Positive Controls: EOC 20 whole cell lysate: sc-364187, PC-12 cell lysate: sc-2250 or C6 whole cell lysate: sc-364373.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



APOBEC4 (S-16): sc-104060. Western blot analysis of APOBEC4 expression in PC-12 (A), EOC 20 (B), BC₃H1 (C) and C6 (D) whole cell lysates and rat epididymis tissue extract (E).

RESEARCH USE

For research use only, not for use in diagnostic procedures.