

CacyBP (C-15): sc-104120

BACKGROUND

CacyBP (calcyclin-binding protein, SIAH-interacting protein) is a 228 amino acid protein encoded by the human gene CACYBP. CacyBP is primarily a nuclear protein that contains one CS domain and one SGS domain. CacyBP is believed to be involved in calcium-dependent ubiquitination and subsequent proteosomal degradation of target proteins. It most likely serves as a molecular bridge in ubiquitin E3 complexes. It also participates in the ubiquitin-mediated degradation of β -catenin. CacyBP is thought to be a potential inhibitor of cell growth and invasion in the gastric cancer cell through its effects on β -catenin protein expression and transcriptional activation of TCF/LEF.

REFERENCES

- Hildebrandt, B., et al. 2007. Differential gene expression in peripheral blood lymphocytes of cancer patients treated with whole body hyperthermia and chemotherapy: a pilot study. *Int. J. Hyperthermia* 22: 625-635.
- Sun, S., et al. 2007. Overexpressed CacyBP/SIP leads to the suppression of growth in renal cell carcinoma. *Biochem. Biophys. Res. Commun.* 356: 864-871.
- Mei, Y., et al. 2007. SIAH-1S, a novel splice variant of SIAH-1 (seven in absentia homolog), counteracts SIAH-1-mediated downregulation of β -catenin. *Oncogene* 26: 6319-6331.
- Herington, J.L., et al. 2007. β -catenin (CTNNB) in the mouse uterus during decidualization and the potential role of two pathways in regulating its degradation. *J. Histochem. Cytochem.* 55: 963-974.
- Liang, J., et al. 2007. Differential expression of calcium-related genes in gastric cancer cells transfected with cellular prion protein. *Biochem. Cell Biol.* 85: 375-383.
- Schneider, G., et al. 2007. CacyBP/SIP interacts with Tubulin in neuroblastoma NB2a cells and induces formation of globular Tubulin assemblies. *Biochim. Biophys. Acta* 1773: 1628-1636.
- Ning, X., et al. 2008. Calcyclin-binding protein inhibits proliferation, tumorigenicity, and invasion of gastric cancer. *Mol. Cancer Res.* 5: 1254-1262.

CHROMOSOMAL LOCATION

Genetic locus: CACYBP (human) mapping to 1q25.1; Cacybp (mouse) mapping to 1 H2.1.

SOURCE

CacyBP (C-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of CacyBP of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-104120 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

CacyBP (C-15) is recommended for detection of CacyBP of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

CacyBP (C-15) is also recommended for detection of CacyBP in additional species, including equine, canine, bovine and avian.

Suitable for use as control antibody for CacyBP siRNA (h): sc-88504, CacyBP siRNA (m): sc-77341, CacyBP shRNA Plasmid (h): sc-88504-SH, CacyBP shRNA Plasmid (m): sc-77341-SH, CacyBP shRNA (h) Lentiviral Particles: sc-88504-V and CacyBP shRNA (m) Lentiviral Particles: sc-77341-V.


Molecular Weight of CacyBP: 26 kDa.

Positive Controls: Hep G2 cell lysate: sc-2227, mouse embryo extract: sc-364239 or CacyBP (m): 293T Lysate: sc-118948.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



CacyBP (C-15): sc-104120. Western blot analysis of CacyBP expression in non-transfected: sc-117752 (A) and mouse CacyBP transfected: sc-118948 (B) 293T whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.