



COX18 (N-14): sc-104164

BACKGROUND

The cytochrome c oxidase (COX) family of proteins function as the final electron donor in the respiratory chain to drive a proton gradient across the inner mitochondrial membrane, ultimately resulting in the production of water. COX18 (cytochrome c oxidase-assembly protein 18), also known as Cytochrome c oxidase assembly protein 18 is a 333 amino acid mitochondrial inner membrane protein that plays a role in the assembly of complex IV of the mitochondrial respiratory chain by catalyzing the insertion of the C-terminal tail of COX2 into the inner membrane. Deficiency of complex IV is a rare metabolic disorder that results in clinically heterogeneous symptoms ranging from isolated myopathy to severe multisystem disease. There are four isoforms of COX18 that are expressed as a result of alternative splicing events.

REFERENCES

1. Sacconi, S., Trevisson, E., Pistollato, F., Baldoiu, M.C., Rezzonico, R., Bourget, I., Desnuelle, C., Tenconi, R., Basso, G., DiMauro, S. and Salviati, L. 2005. hCOX18 and hCOX19: two human genes involved in cytochrome c oxidase assembly. *Biochem. Biophys. Res. Commun.* 337: 832-839.
2. Gaisne, M. and Bonnefoy, N. 2006. The COX18 gene, involved in mitochondrial biogenesis, is functionally conserved and tightly regulated in humans and fission yeast. *FEMS Yeast Res.* 6: 869-882.
3. Online Mendelian Inheritance in Man, OMIM™. 2006. Johns Hopkins University, Baltimore, MD. MIM Number: 610428. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. van Bloois, E., Koningstein, G., Bauerschmitt, H., Herrmann, J.M. and Luirink, J. 2007. *Saccharomyces cerevisiae* COX18 complements the essential Sec-independent function of *Escherichia coli* YidC. *FEBS J.* 274: 5704-5713.
5. Jia, L., Dienhart, M.K. and Stuart, R.A. 2007. Oxa1 directly interacts with Atp9 and mediates its assembly into the mitochondrial F1Fo-ATP synthase complex. *Mol. Biol. Cell.* 18: 1897-1908.
6. Fiumera, H.L., Broadley, S.A. and Fox, T.D. 2007. Translocation of mitochondrially synthesized Cox2 domains from the matrix to the intermembrane space. *Mol. Cell. Biol.* 27: 4664-4673.
7. Bonnefoy, N., Fiumera, H.L., Dujardin, G. and Fox, T.D. 2009. Roles of Oxa1-related inner-membrane translocases in assembly of respiratory chain complexes. *Biochim. Biophys. Acta* 1793: 60-70.
8. Sacconi, S., Salviati, L. and Trevisson, E. 2009. Mutation analysis of COX18 in 29 patients with isolated cytochrome c oxidase deficiency. *J. Hum. Genet.* 54: 419-421.
9. Online Mendelian Inheritance in Man, OMIM™. 2009. Johns Hopkins University, Baltimore, MD. MIM Number: 220110. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: COX18 (human) mapping to 4q13.3.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

SOURCE

COX18 (N-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of COX18 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-104164 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

COX18 (N-14) is recommended for detection of COX18 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other COX family members.

Suitable for use as control antibody for COX18 siRNA (h): sc-89246, COX18 shRNA Plasmid (h): sc-89246-SH and COX18 shRNA (h) Lentiviral Particles: sc-89246-V.

Molecular Weight of COX18: 37 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.