

KLF14 (C-19): sc-104345

BACKGROUND

Krüppel-like factors (KLFs) comprise a family of evolutionarily conserved zinc finger-containing transcription factors with diverse regulatory functions in cell growth, proliferation, differentiation and embryogenesis. Individual members of the Sp1-like/KLF family can function either as activators or repressors, depending on which promoter they bind and which coregulators they interact with. KLF14 (Krüppel-like factor 14), also known as BTEB5 (basic transcription element-binding protein 5), is a 323 amino acid protein that localizes to the nucleus and contains three C₂H₂-type zinc fingers, suggesting a role in transcriptional regulation. The gene encoding KLF14 maps to human chromosome 7, which houses over 1,000 genes and comprises nearly 5% of the human genome. Defects in some of the genes localized to chromosome 7 have been linked to osteogenesis imperfecta, Williams-Beuren syndrome, Pendred syndrome, lissencephaly, citrullinemia and Shwachman-Diamond syndrome.

REFERENCES

- Dang, D.T., Pevsner, J. and Yang, V.W. 2000. The biology of the mammalian Krüppel-like family of transcription factors. *Int. J. Biochem. Cell Biol.* 32: 1103-1121.
- Black, A.R., Black, J.D. and Azizkhan-Clifford, J. 2001. Sp1 and Krüppel-like factor family of transcription factors in cell growth regulation and cancer. *J. Cell. Physiol.* 188: 143-160.
- Kaczynski, J., Cook, T. and Urrutia, R. 2003. Sp1- and Krüppel-like transcription factors. *Genome Biol.* 4: 206.
- Suske, G., Bruford, E. and Philipsen, S. 2005. Mammalian SP/KLF transcription factors: bring in the family. *Genomics* 85: 551-556.
- Parker-Katirae, L., Carson, A.R., Yamada, T., Arnaud, P., Feil, R., Abu-Amero, S.N., Moore, G.E., Kaneda, M., Perry, G.H., Stone, A.C., Lee, C., Meguro-Horike, M., Sasaki, H., Kobayashi, K., Nakabayashi, K. and Scherer, S.W. 2007. Identification of the imprinted KLF14 transcription factor undergoing human-specific accelerated evolution. *PLoS Genet.* 3: e65.
- Online Mendelian Inheritance in Man, OMIM[™]. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 609393. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Truty, M.J., Lomber, G., Fernandez-Zapico, M.E. and Urrutia, R. 2009. Silencing of the transforming growth factor- β (TGF β) receptor II by Krüppel-like factor 14 underscores the importance of a negative feedback mechanism in TGF β signaling. *J. Biol. Chem.* 284: 6291-6300.

CHROMOSOMAL LOCATION

Genetic locus: KLF14 (human) mapping to 7q32.2.

SOURCE

KLF14 (C-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of KLF14 of human origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-104345 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

KLF14 (C-19) is recommended for detection of KLF14 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for KLF14 siRNA (h): sc-89527, KLF14 shRNA Plasmid (h): sc-89527-SH and KLF14 shRNA (h) Lentiviral Particles: sc-89527-V.

Molecular Weight of KLF14: 33 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.