

LIN-9 (T-17): sc-104353

BACKGROUND

LIN-9, also known as TGS, BARA or TGS1, is a 542 amino acid protein that localizes to the nucleoplasm and is a mammalian homolog of the *C. elegans* Lin-9 protein. Expressed in testis and thymus, LIN-9 functions as a component of the DREAM complex (also known as the LINC complex), which is comprised of several proteins, all of which work in concert to repress cell cycle-dependent genes. LIN-9 specifically acts as a tumor suppressor that associates with Rb and inhibits DNA synthesis, possibly also controlling the expression of genes that are required for the G₁/S cell cycle transition. Three isoforms of LIN-9 exist due to alternative splicing events. The gene encoding LIN-9 maps to human chromosome 1, which spans 260 million base pairs, contains over 3,000 genes and comprises nearly 8% of the human genome.

REFERENCES

- Korenjak, M., et al. 2004. Native E2F/RBF complexes contain Myb-interacting proteins and repress transcription of developmentally controlled E2F target genes. *Cell* 119: 181-193.
- Gagrica, S., et al. 2004. Inhibition of oncogenic transformation by mammalian LIN-9, a pRb-associated protein. *EMBO J.* 23: 4627-4638.
- Online Mendelian Inheritance in Man, OMIM™. 2005. Johns Hopkins University, Baltimore, MD. MIM Number: 609375. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Sandoval, R., et al. 2006. A mutant allele of BARA/LIN-9 rescues the Cdk4^{-/-} phenotype by releasing the repression on E2F-regulated genes. *Exp. Cell Res.* 312: 2465-2475.
- Osterloh, L., et al. 2007. The human synMuv-like protein LIN-9 is required for transcription of G₂/M genes and for entry into mitosis. *EMBO J.* 26: 144-157.
- Pilkinton, M., et al. 2007. Mip/LIN-9 regulates the expression of B-Myb and the induction of cyclin A, cyclin B, and Cdk1. *J. Biol. Chem.* 282: 168-175.
- Pilkinton, M., et al. 2007. Mammalian Mip/LIN-9 interacts with either the p107, p130/E2F4 repressor complex or B-Myb in a cell cycle-phase-dependent context distinct from the *Drosophila* dREAM complex. *Oncogene* 26: 7535-7543.

CHROMOSOMAL LOCATION

Genetic locus: LIN9 (human) mapping to 1q42.12; Lin9 (mouse) mapping to 1 H4.

SOURCE

LIN-9 (T-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of LIN-9 of human origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-104353 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-104353 X, 200 µg/0.1 ml.

APPLICATIONS

LIN-9 (T-17) is recommended for detection of LIN-9 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other LIN family members.

LIN-9 (T-17) is also recommended for detection of LIN-9 in additional species, including equine, canine and avian.

Suitable for use as control antibody for LIN-9 siRNA (h): sc-88786, LIN-9 siRNA (m): sc-105615, LIN-9 shRNA Plasmid (h): sc-88786-SH, LIN-9 shRNA Plasmid (m): sc-105615-SH, LIN-9 shRNA (h) Lentiviral Particles: sc-88786-V and LIN-9 shRNA (m) Lentiviral Particles: sc-105615-V.

LIN-9 (T-17) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Molecular Weight of LIN-9 isoforms 1/2/3: 62/64/58 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



MONOS
Satisfaction
Guaranteed

Try **LIN-9 (C-10): sc-398234** or **LIN-9 (Mab#1): sc-130571**, our highly recommended monoclonal alternatives to LIN-9 (T-17).