



Pdcd-7 (K-14): sc-104585

BACKGROUND

PD-1 (programmed cell death-1 protein) is a type I transmembrane receptor and a member of the immunoglobulin gene superfamily. Expression of PD-1 is detected in mouse thymus, and it is induced in stimulated B and T cell lines, where it may play a role in the negative regulation of various immune responses. PD-2 (also known as PDCD2, PDL2 or B7DC) is highly expressed in placenta, heart, pancreas, lung and liver. PD-2 is also expressed in spleen, lymph nodes and thymus. Pdcd-4 is a nuclear protein localized to the nucleus of proliferating cells. ALG-2 (apoptosis-linked gene 2), also designated programmed cell death protein 6, is a Ca²⁺-binding protein that participates in regulatory events occurring late in the apoptotic program and where several death signals converge. Pdcd-7, also designated programmed cell death protein 7, is a 485 amino acid protein that promotes apoptosis when over-expressed. Human Pdcd-7 and the mouse Pdcd-7 protein, designated as Es18, share 95% homology in the overlapping region.

REFERENCES

1. Ishida, Y., Agata, Y., Shibahara, K. and Honjo, T. 1992. Induced expression of PD-1, a novel member of the immunoglobulin gene superfamily, upon programmed cell death. *EMBO J.* 11: 3887-3895.
2. Agata, Y., Kawasaki, A., Nishimura, H., Ishida, Y., Tsubata, T., Yagita, H. and Honjo, T. 1996. Expression of the PD-1 antigen on the surface of stimulated mouse T and B lymphocytes. *Int. Immunol.* 8: 765-772.
3. Park, E.J., Kim, J.H., Seong, R.H., Kim, C.G., Park, S.D. and Hong, S.H. 1999. Characterization of a novel mouse cDNA, ES18, involved in apoptotic cell death of T cells. *Nucleic Acids Res.* 27: 1524-1530.
4. Li, W., Jin, K., Nagayama, T., He, X., Chang, J., Minami, M., Graham, S.H. and Simon, R.P. and Greenberg, D.A. 2000. Increased expression of apoptosis-linked gene 2 (ALG2) in the rat brain after temporary focal cerebral ischemia. *Neuroscience* 96: 161-168.
5. Latchman, Y., Wood, C.R., Chernova, T., Chaudhary, D., Borde, M., Chernova, I., Iwai, Y., Long, A.J., Brown, J.A., Nunes, R., Greenfield, E.A., Bourque, K., Boussiotis, V.A., Carter, L.L., Carreno, B.M., Malenkovich, N., Nishimura, H., Okazaki, T., Honjo, T., Sharpe, A.H. and Freeman, G.J. 2001. PD-L2 is a second ligand for PD-1 and inhibits T cell activation. *Nat. Immunol.* 2: 261-268.
6. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 608138. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
7. Liu, Z. and Aune, T.M. 2006. Deregulated stress system in non-obese diabetic lymphocyte. *Genes Immun.* 7: 352-358.

CHROMOSOMAL LOCATION

Genetic locus: PDCD7 (human) mapping to 15q22.31; Pdcd7 (mouse) mapping to 9 C.

SOURCE

Pdcd-7 (K-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Pdcd-7 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-104585 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Pdcd-7 (K-14) is recommended for detection of Pdcd-7 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Pdcd-7 siRNA (h): sc-90046, Pdcd-7 siRNA (m): sc-72132, Pdcd-7 shRNA Plasmid (h): sc-90046-SH, Pdcd-7 shRNA Plasmid (m): sc-72132-SH, Pdcd-7 shRNA (h) Lentiviral Particles: sc-90046-V and Pdcd-7 shRNA (m) Lentiviral Particles: sc-72132-V.

Molecular Weight of Pdcd-7: 55 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200 or mouse testis tissue extract.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.