



Sp8 (N-14): sc-104662

BACKGROUND

Sp8, also known as BTD, is a 508 amino acid nuclear protein that contains three C₂H₂-type zinc fingers. Existing as multiple alternatively spliced isoforms, Sp8 plays a crucial role in limb outgrowth and neuropore closure and is thought to mediate apical ectodermal ridge (AER) formation. The gene encoding Sp8 maps to human chromosome 7, which houses over 1,000 genes and comprises nearly 5% of the human genome. Defects in some of the genes localized to chromosome 7 have been linked to Osteogenesis imperfecta, Pendred syndrome, Lissencephaly, Citrullinemia and Shwachman-Diamond syndrome. The deletion of a portion of the q arm of chromosome 7 is associated with Williams-Beuren syndrome, a condition characterized by mild mental retardation, an unusual comfort and friendliness with strangers and an elfin appearance. Deletions of portions of the q arm of chromosome 7 are also seen in a number of myeloid disorders, including cases of acute myelogenous leukemia and myelodysplasia.

REFERENCES

1. Treichel, D., Schöck, F., Jäckle, H., Gruss, P. and Mansouri, A. 2003. MBtd is required to maintain signaling during murine limb development. *Genes Dev.* 17: 2630-2635.
2. Bell, S.M., Schreiner, C.M., Waclaw, R.R., Campbell, K., Potter, S.S. and Scott, W.J. 2003. Sp8 is crucial for limb outgrowth and neuropore closure. *Proc. Natl. Acad. Sci. USA* 100: 12195-12200.
3. Milona, M.A., Gough, J.E. and Edgar, A.J. 2004. Genomic structure and cloning of two transcript isoforms of human Sp8. *BMC Genomics* 5: 86.
4. Online Mendelian Inheritance in Man, OMIM™. 2005. Johns Hopkins University, Baltimore, MD. MIM Number: 608306. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Eckert, M.A., Galaburda, A.M., Mills, D.L., Bellugi, U., Korenberg, J.R. and Reiss, A.L. 2006. The neurobiology of Williams syndrome: cascading influences of visual system impairment? *Cell. Mol. Life Sci.* 63: 1867-1875.
6. Osborne, L.R., Joseph-George, A.M. and Scherer, S.W. 2006. Williams-Beuren syndrome diagnosis using fluorescence *in situ* hybridization. *Methods Mol. Med.* 126:113-128.
7. Reiner, O., Sapoznik, S. and Sapir, T. 2006. Lissencephaly 1 linking to multiple diseases: mental retardation, neurodegeneration, schizophrenia, male sterility, and more. *Neuromolecular Med.* 8: 547-565.
8. Shimamura, A. 2006. Shwachman-Diamond syndrome. *Semin. Hematol.* 43: 178-188.
9. Brezinová, J., Zemanová, Z., Ransdorfová, S., Pavlistová, L., Babická, L., Housková, L., Melicherciková, J., Sisková, M., Cermák, J. and Michalová, K. 2007. Structural aberrations of chromosome 7 revealed by a combination of molecular cytogenetic techniques in myeloid malignancies. *Cancer Genet. Cytogenet.* 173: 10-16.

CHROMOSOMAL LOCATION

Genetic locus: SP8 (human) mapping to 7p15.3; Sp8 (mouse) mapping to 12 F2.

SOURCE

Sp8 (N-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Sp8 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-104662 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Sp8 (N-14) is recommended for detection of Sp8 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for SP8 siRNA (h): sc-89539, Sp8 siRNA (m): sc-153692, SP8 shRNA Plasmid (h): sc-89539-SH, Sp8 shRNA Plasmid (m): sc-153692-SH, SP8 shRNA (h) Lentiviral Particles: sc-89539-V and Sp8 shRNA (m) Lentiviral Particles: sc-153692-V.

Molecular Weight of Sp8: 51 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.