

TMED4 (Q-12): sc-104698

BACKGROUND

TMED4 (transmembrane emp24 domain-containing protein 4), also known as ERS25 (endoplasmic reticulum stress-response protein 25) and putative NF κ B-activating protein 156, is a 225 amino acid protein that is a member of the EMP24/GP25L family. Like most members of this family, TMED4 is a single-pass type I membrane protein containing one GOLD domain. The GOLD (Golgi dynamics) domain is a region of about 90 to 150 amino acids that mediates protein-protein interactions. The GOLD domain interacts with lipid, sterol or fatty acid-domains as well as with the RUN domain, which interacts with cytoskeletal filaments, of membrane proteins. Localized to the endoplasmic reticulum, TMED4 is induced by ER-specific stress, heat shock and oxidative stress. Knockdown of TMED4 mRNA results in a significant reduction in apoptosis as well as a reduction in reactive oxidative species. There are three isoforms of TMED4 that are produced as a result of alternative splicing events.

REFERENCES

- Dominguez, M., et al. 1998. GP25L/EMP24/p24 protein family members of the *cis*-Golgi network bind both COP I and II coatomer. *J. Cell Biol.* 140: 751-765.
- Anantharaman, V. and Aravind, L. 2002. The GOLD domain, a novel protein module involved in Golgi function and secretion. *Genome Biol.* 3: research0023.
- Matsuda, A., et al. 2003. Large-scale identification and characterization of human genes that activate NF κ B and MAPK signaling pathways. *Oncogene* 22: 3307-3318.
- Nakanishi, K., et al. 2004. Localization and function in endoplasmic reticulum stress tolerance of ERdj3, a new member of Hsp40 family protein. *Cell Stress Chaperones* 9: 253-264.
- Ishiyama, T., et al. 2007. OCIA domain containing 2 is highly expressed in adenocarcinoma mixed subtype with bronchioloalveolar carcinoma component and is associated with better prognosis. *Cancer Sci.* 98: 50-57.
- Hwang, S.O., et al. 2008. Novel oxidative stress-responsive gene ERS25 functions as a regulator of the heat-shock and cell death response. *J. Biol. Chem.* 283: 13063-13069.
- Online Mendelian Inheritance in Man, OMIM™. 2008. Johns Hopkins University, Baltimore, MD. MIM Number: 612038. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Chen, R., et al. 2009. Glycoproteomics analysis of human liver tissue by combination of multiple enzyme digestion and hydrazide chemistry. *J. Proteome Res.* 8: 651-661.

CHROMOSOMAL LOCATION

Genetic locus: TMED4 (human) mapping to 7p13; Tmed4 (mouse) mapping to 11 A1.

SOURCE

TMED4 (Q-12) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of TMED4 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-104698 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

TMED4 (Q-12) is recommended for detection of TMED4 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

TMED4 (Q-12) is also recommended for detection of TMED4 in additional species, including porcine.

Suitable for use as control antibody for TMED4 siRNA (h): sc-89384, TMED4 siRNA (m): sc-106619, TMED4 shRNA Plasmid (h): sc-89384-SH, TMED4 shRNA Plasmid (m): sc-106619-SH, TMED4 shRNA (h) Lentiviral Particles: sc-89384-V and TMED4 shRNA (m) Lentiviral Particles: sc-106619-V.

Molecular Weight of TMED4 isoforms: 26/24/21 kDa.

Positive Controls: EOC 20 whole cell lysate: sc-364187.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.