MSE55 (S-20): sc-10491



The Power to Question

BACKGROUND

MSE55 also known as Borg5 or binder of Rho GTPases 5 is a Cdc42 effector protein that induces long cellular extensions in fibroblasts. MSE55 also increases and mediates actin cytoskeleton reorganization at the plasma membrane. MSE55 is a nonkinase CRIB (Cdc42/Rac interactive-binding) domain-containing molecule. An intact CRIB domain is required for the GTP-dependent binding of MSE55 to Cdc42. MSE55 is expressed in a tissue-specific manner in endothelial and bone marrow stromal cells. MSE55 may have a functional role in hematopoiesis or as a negative regulator of Rho GTPase signaling.

REFERENCES

- Bahou, W.F., Campbell, A.D. and Wicha, M.S. 1992. cDNA cloning and molecular characterization of MSE55, a novel human serum constituent protein that displays bone marrow stromal/endothelial cell-specific expression. J. Biol. Chem. 267: 13986-13992.
- Burbelo, P.D., Drechsel, D. and Hall, A. 1995. A conserved binding motif defines numerous candidate target proteins for both Cdc42 and Rac GTPases. J. Biol. Chem. 270: 29071-29074.
- Neudauer, C.L., Joberty, G., Tatsis, N. and Macara, I.G. 1998. Distinct cellular effects and interactions of the Rho-family GTPase TC10. Curr. Biol. 8: 1151-1160.
- 4. Burbelo, P.D., Snow, D.M., Bahou, W. and Spiegel, S. 1999. MSE55, a Cdc42 effector protein, induces long cellular extensions in fibroblasts. Proc. Natl. Acad. Sci. USA 96: 9083-9088.
- 5. Joberty, G., Perlungher, R.R. and Macara, I.G. 1999. The Borgs, a new family of Cdc42 and TC10 GTPase-interacting proteins. Mol. Cell. Biol. 19: 6585-6597.

CHROMOSOMAL LOCATION

Genetic locus: CDC42EP1 (human) mapping to 22q13.1.

SOURCE

MSE55 (S-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of MSE55 of human origin.

PRODUCT

Each vial contains 200 μg lgG in 1.0 ml of PBS with <0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-10491 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

MSE55 (S-20) is recommended for detection of MSE55 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

MSE55 (S-20) is also recommended for detection of MSE55 in additional species, including canine, bovine and porcine.

Suitable for use as control antibody for MSE55 siRNA (h): sc-41802, MSE55 shRNA Plasmid (h): sc-41802-SH and MSE55 shRNA (h) Lentiviral Particles: sc-41802-V.


Molecular Weight of MSE55: 55 kDa.

Positive Controls: ECV304 cell lysate: sc-2269 or MSE55 (h3): 293T Lysate: sc-175134.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



MSE55 (S-20): sc-10491. Western blot analysis of MSE55 expression in non-transfected 293T: sc-117752 (**A**), human MSE55 transfected 293T: sc-175134 (**B**) and ECV304 (**C**) whole cell Ivsates.

RESEARCH USE

For research use only, not for use in diagnostic procedures

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3800 fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com