

RFX5 (E-17): sc-10667

BACKGROUND

EP is a regulatory enhancer element found in several promoters on viral genes, and similar sites are also present in cellular genes, including the MIF-1 binding site (MIE) of the human c-Myc gene, the X box of MHC class II promoters and a binding site in the proliferating cell nuclear antigen promoter. The EP sites present in the X box of MHC class II promoters are distinctly nonpalindromic sequences that contain only a single EP-homologous half-site. The EP element is bound by an ubiquitous nuclear protein complex that consists of homo- and heterodimers involving the RFX1, RFX2 and RFX3 proteins. The RFX proteins represent an essential class II transcription factor family that shares several conserved regions, including the centrally located DNA-binding domain (DBD) and the D region found in the C-terminal part of these proteins which facilitates dimerization. RFX complexes can activate the enhancer elements of several HBV genes and also promote the induction of MHC class II genes in response to interferon- γ stimulation. Two additional subunits, RFX5, RFX-B/Ank, are also involved in the RFX complexes, yet they bind additional elements in the X1 half of the X box.

REFERENCES

1. Dikstein, R., et al. 1990. Functional organization of the hepatitis B virus enhancer. *Mol. Cell. Biol.* 10: 3682-3689.
2. Fontes, J.D., et al. 1997. Assembly of functional regulatory complexes on MHC class II promoters *in vivo*. *J. Mol. Biol.* 270: 336-345.
3. Katan, Y., et al. 1997. The transcriptional activation and repression domains of RFX1, a context-dependent regulator, can mutually neutralize their activities. *Nucleic Acids Res.* 25: 3621-3628.
4. Masternak, K., et al. 1998. A gene encoding a novel RFX-associated transactivator is mutated in the majority of MHC class II deficiency patients. *Nat. Genet.* 20: 273-277.
5. Katan-Khaykovich, Y., et al. 1998. RFX1, a single DNA-binding protein with a split dimerization domain, generates alternative complexes. *J. Biol. Chem.* 273: 24504-24512.

CHROMOSOMAL LOCATION

Genetic locus: RFX5 (human) mapping to 1q21.3.

SOURCE

RFX5 (E-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of RFX5 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-10667 X, 200 μ g/0.1 ml.

Blocking peptide available for competition studies, sc-10667 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

RFX5 (E-17) is recommended for detection of RFX5 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Suitable for use as control antibody for RFX5 siRNA (h): sc-37747, RFX5 shRNA Plasmid (h): sc-37747-SH and RFX5 shRNA (h) Lentiviral Particles: sc-37747-V.

RFX5 (E-17) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of RFX5: 75 kDa.

Positive Controls: RFX5 (h): 293 Lysate: sc-111879 or Jurkat nuclear extract: sc-2132.

DATA



RFX5 (E-17): sc-10667. Western blot analysis of RFX5 expression in non-transfected: sc-110760 (A) and human RFX5 transfected: sc-111879 (B) 293 whole cell lysates.



RFX5 (E-17): sc-10667. Western blot analysis of RFX5 expression in Jurkat nuclear extract.

SELECT PRODUCT CITATIONS

1. Wolfe, S.A., et al. 2006. Transcription factor RFX2 is abundant in rat testis and enriched in nuclei of primary spermatocytes where it appears to be required for transcription of the testis-specific histone H1t gene. *J. Cell. Biochem.* 99: 735-746.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfaction
Guaranteed

Try **RFX5 (C-3): sc-271756** or **RFX5 (A-10): sc-271757**, our highly recommended monoclonal alternatives to RFX5 (E-17).