

TIP120B (T-20): sc-10677

BACKGROUND

TATA-binding protein (TBP) forms complexes with various nuclear proteins and is a target for various transcriptional regulators, such as TIP120. The two members of the TIP120 family of proteins, TIP120A and TIP120B, are TBP-Interacting Proteins that function as global activators in transcriptional regulation. TIP120A is a ubiquitously expressed 120 kDa protein isolated from rat liver nuclear extracts, originally named TIP120. TIP120B is a TIP120A-like protein that is expressed specifically in muscle tissues. TIP120A binds directly to TBP and a particular subunit of RNA polymerases (RNAP) to facilitate specific integration of RNAP II into the preinitiation complex (PIC). In addition to being a transcription factor of TBP, the chaperone-like activity toward the RNA polymerases demonstrates that TIP120 regulates the amplification of multiple gene expression.

REFERENCES

1. Zawel, L., et al. 1992. Advances in RNA polymerase II transcription. *Curr. Opin. Cell. Biol.* 4: 488-495.
2. Conaway, R.C., et al. 1993. General initiation factors for RNA polymerase II. *Annu. Rev. Biochem.* 62: 161-190.
3. Yogosawa, S., et al. 1996. Molecular cloning of a novel 120-kDa TBP-interacting protein. *Biochem. Biophys. Res. Commun.* 229: 612-617.
4. Roeder, R.G. 1996. The role of general initiation factors in transcription by RNA polymerase II. *Trends Biochem. Sci.* 21: 327-335.
5. Aoki, T., et al. 1999. TIP120B: a novel TIP120-family protein that is expressed specifically in muscle tissues. *Biochem. Biophys. Res. Commun.* 261: 911-916.
6. Makino, Y., et al. 1999. TATA-Binding Protein-Interacting Protein 120, TIP120, Stimulates Three Classes of Eukaryotic Transcription via a Unique Mechanism. *Mol. Cell. Biol.* 19: 7951-7960.

SOURCE

TIP120B (T-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of TIP120B of rat origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-10677 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-10677 X, 200 µg/0.1 ml.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

TIP120B (T-20) is recommended for detection of TIP120B, TIP120B short form and TIP120B alternatively spliced form of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1–2 µg per 100–500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for TIP120B siRNA (m): sc-36682.

TIP120B (T-20) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of TIP120B: 151 kDa.

Positive Controls: rat skeletal muscle extract.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



TIP120B (T-20): sc-10677. Western blot analysis of TIP120B expression in rat skeletal muscle extract.

RESEARCH USE

For research use only, not for use in diagnostic procedures.