

MRP-S2 (S-12): sc-107015

BACKGROUND

Mitochondrial ribosomes consist of a large 39S subunit and a small 28S subunit, both of which are comprised of multiple mitochondrial ribosomal proteins (MRPs) that are encoded by nuclear genes and are essential for protein synthesis within mitochondria. MRP-S2 (mitochondrial ribosomal protein S2), also known as S2mt or CGI-91, is a 296 amino acid mitochondrial protein that exists as a component of the 28S subunit and may play a role in mitochondrial protein synthesis. The gene encoding MRP-S2 maps to human chromosome 9, which houses over 900 genes and comprises nearly 4% of the human genome. Hereditary hemorrhagic telangiectasia, which is characterized by harmful vascular defects, and Familial dysautonomia, are both associated with chromosome 9. Notably, chromosome 9 encompasses the largest interferon family gene cluster.

REFERENCES

1. Kenmochi, N., Suzuki, T., Uechi, T., Magoori, M., Kuniba, M., Higa, S., Watanabe, K. and Tanaka, T. 2001. The human mitochondrial ribosomal protein genes: mapping of 54 genes to the chromosomes and implications for human disorders. *Genomics* 77: 65-70.
2. Cavdar Koc, E., Burkhart, W., Blackburn, K., Moseley, A. and Spremulli, L.L. 2001. The small subunit of the mammalian mitochondrial ribosome. Identification of the full complement of ribosomal proteins present. *J. Biol. Chem.* 276: 19363-19374.
3. Koc, E.C., Burkhart, W., Blackburn, K., Moyer, M.B., Schlatzer, D.M., Moseley, A. and Spremulli, L.L. 2001. The large subunit of the mammalian mitochondrial ribosome. Analysis of the complement of ribosomal proteins present. *J. Biol. Chem.* 276: 43958-43969.
4. Zhang, Z. and Gerstein, M. 2003. Identification and characterization of over 100 mitochondrial ribosomal protein pseudogenes in the human genome. *Genomics* 81: 468-480.
5. O'Brien, T.W. 2003. Properties of human mitochondrial ribosomes. *IUBMB Life* 55: 505-513.
6. Sylvester, J.E., Fischel-Ghodsian, N., Mougey, E.B. and O'Brien, T.W. 2004. Mitochondrial ribosomal proteins: candidate genes for mitochondrial disease. *Genet. Med.* 6: 73-80.
7. Online Mendelian Inheritance in Man, OMIM[™]. 2008. Johns Hopkins University, Baltimore, MD. MIM Number: 611971. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: MRPS2 (human) mapping to 9q34.3; Mrps2 (mouse) mapping to 2 A3.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

SOURCE

MRP-S2 (S-12) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of MRP-S2 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-107015 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

MRP-S2 (S-12) is recommended for detection of MRP-S2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

MRP-S2 (S-12) is also recommended for detection of MRP-S2 in additional species, including canine and bovine.

Suitable for use as control antibody for MRP-S2 siRNA (h): sc-92589, MRP-S2 siRNA (m): sc-149621, MRP-S2 shRNA Plasmid (h): sc-92589-SH, MRP-S2 shRNA Plasmid (m): sc-149621-SH, MRP-S2 shRNA (h) Lentiviral Particles: sc-92589-V and MRP-S2 shRNA (m) Lentiviral Particles: sc-149621-V.

Molecular Weight of MRP-S2: 33 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.