

RNF130 (S-15): sc-107089

BACKGROUND

The RING-type zinc finger motif is present in a number of viral and eukaryotic proteins and is made of a conserved cysteine-rich domain that is able to bind two zinc atoms. Proteins that contain this conserved domain are generally involved in the ubiquitination pathway of protein degradation. RNF130 (ring finger protein 130), also known as GP, G1RZFP (G₁-related zinc finger protein) or GOLIATH, is a 419 amino acid single-pass type I membrane protein that shares similarity with a *Drosophila* zinc-finger protein found in mesoderm known as γ 1. RNF130 contains one PA (protease associated) domain and a single RING-type zinc finger. Implicated in the regulation of growth factor withdrawal-induced apoptosis of myeloid precursor cells, RNF130 is encoded by a gene located on human chromosome 5q35.3 and mouse chromosome 11 B1.3.

REFERENCES

1. Borden, K.L. and Freemont, P.S. 1996. The RING finger domain: a recent example of a sequence-structure family. *Curr. Opin. Struct. Biol.* 6: 395-401.
2. Saurin, A.J., Borden, K.L., Boddy, M.N. and Freemont, P.S. 1996. Does this have a familiar RING? *Trends Biochem. Sci.* 21: 208-214.
3. Lorick, K.L., Jensen, J.P., Fang, S., Ong, A.M., Hatakeyama, S. and Weissman, A.M. 1999. RING fingers mediate ubiquitin-conjugating enzyme (E2)-dependent ubiquitination. *Proc. Natl. Acad. Sci. USA* 96: 11364-11369.

CHROMOSOMAL LOCATION

Genetic locus: RNF130 (human) mapping to 5q35.3; Rnf130 (mouse) mapping to 11 B1.3.

SOURCE

RNF130 (S-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping within a C-terminal cytoplasmic domain of RNF130 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-107089 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-107089 X, 200 μ g/0.1 ml.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

RNF130 (S-15) is recommended for detection of RNF130 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

RNF130 (S-15) is also recommended for detection of RNF130 in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for RNF130 siRNA (h): sc-91644, RNF130 siRNA (m): sc-153010, RNF130 shRNA Plasmid (h): sc-91644-SH, RNF130 shRNA Plasmid (m): sc-153010-SH, RNF130 shRNA (h) Lentiviral Particles: sc-91644-V and RNF130 shRNA (m) Lentiviral Particles: sc-153010-V.

RNF130 (S-15) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of RNF130: 46 kDa.

Positive Controls: RNF130 (m): 293T Lysate: sc-125933.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



RNF130 (S-15): sc-107089. Western blot analysis of RNF130 expression in non-transfected: sc-117752 (A) and mouse RNF130 transfected: sc-125933 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.