

ERp5 (N-16): sc-107533

BACKGROUND

Endoplasmic reticulum proteins (ERPs) are widely expressed proteins that localize to the ER and may act as proteases, protein disulfide isomerases, thiol-disulfide oxidases or phospholipases. ERp5, also known as PDIA6 (protein disulfide isomerase family A, member 6) or TXNDC7, is a 440 amino acid protein that contains 2 thioredoxin domains and belongs to the protein disulfide isomerase family. Localized to the melanosome, as well as to the lumen of the endoplasmic reticulum, ERp5 functions to catalyze the rearrangement of disulfide bonds in a variety of different proteins. Via its catalytic activity, ERp5 is able to reduce the disulfide bond that binds MICA to tumor cells, thereby releasing MICA and reducing the rate of tumor expansion. Multiple isoforms of ERp5 exist due to alternative splicing events.

REFERENCES

1. Chaudhuri, M.M., et al. 1992. The gene for a novel protein, a member of the protein disulphide isomerase/form I phosphoinositide-specific phospholipase C family, is amplified in hydroxyurea-resistant cells. *Biochem. J.* 281: 645-650.
2. Hayano, T., et al. 1995. Cloning and sequencing of the cDNA encoding human p5. *Gene* 164: 377-378.
3. Hoshijima, K., et al. 2002. A protein disulfide isomerase expressed in the embryonic midline is required for left/right asymmetries. *Genes Dev.* 16: 2518-2529.
4. Kikuchi, M., et al. 2002. Functional analysis of human p5, a protein disulfide isomerase homologue. *J. Biochem.* 132: 451-455.
5. Jordan, P.A., et al. 2005. A role for the thiol isomerase protein ERp5 in platelet function. *Blood* 105: 1500-1507.

CHROMOSOMAL LOCATION

Genetic locus: PDIA6 (human) mapping to 2p25.1; Pdia6 (mouse) mapping to 12 A1.1.

SOURCE

ERp5 (N-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of ERp5 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-107533 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

ERp5 (N-16) is recommended for detection of ERp5 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other ERp family members.

ERp5 (N-16) is also recommended for detection of ERp5 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for ERp5 siRNA (h): sc-94493, ERp5 siRNA (m): sc-144938, ERp5 shRNA Plasmid (h): sc-94493-SH, ERp5 shRNA Plasmid (m): sc-144938-SH, ERp5 shRNA (h) Lentiviral Particles: sc-94493-V and ERp5 shRNA (m) Lentiviral Particles: sc-144938-V.


Molecular Weight of ERp5: 48 kDa.

Positive Controls: NAMALWA cell lysate: sc-2234, Hep G2 cell lysate: sc-2227 or Raji whole cell lysate: sc-364236.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



ERp5 (N-16): sc-107533. Western blot analysis of ERp5 expression in NAMALWA (A), Raji (B) and Hep G2 (C) whole cell lysates.



ERp5 (N-16): sc-107533. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic localization.

SELECT PRODUCT CITATIONS

1. Huergo-Zapico, L., et al. 2012. Expression of ERp5 and GRP78 on the membrane of chronic lymphocytic leukemia cells: association with soluble MICA shedding. *Cancer Immunol. Immunother.* 61: 1201-1210.
2. Tufo, G., et al. 2014. The protein disulfide isomerases PDIA4 and PDIA6 mediate resistance to cisplatin-induced cell death in lung adenocarcinoma. *Cell Death Differ.* 21: 685-695.