

GalNAc-T4 (N-12): sc-107552

BACKGROUND

The UDP-N-acetyl- α -D-galactosamine:polypeptide N-acetylgalactosaminyltransferase (GalNAc-T) family of enzymes are substrate-specific proteins that catalyze the transfer of GalNAc (N-acetylgalactosamine) to serine and threonine residues onto various proteins, thereby initiating mucin-type O-linked glycosylation in the Golgi apparatus. GalNAc-T4 (Polypeptide N-acetylgalactosaminyltransferase 4), also known as UDP-GalNAc:polypeptide N-acetylgalactosaminyltransferase 4, is a 578 amino acid protein that displays high enzymatic activity toward Muc7, Muc2 and EA2. GalNAc-T4 also glycosylates PSGL-1 on threonine 57. Its N-terminal domain is involved in substrate binding and manganese coordination, while the C-terminal domain is involved in UDP-Gal binding and catalytic reaction. Though ubiquitously expressed, GalNAc-T4 is found at highest levels in mucosal cells.

REFERENCES

- Bennett, E.P., et al. 1996. cDNA cloning and expression of a novel human UDP-N-acetyl- α -D-galactosamine. Polypeptide N-acetylgalactosaminyltransferase, GalNAc-t3. *J. Biol. Chem.* 271: 17006-17012.
- Bennett, E.P., et al. 1998. Cloning of a human UDP-N-acetyl- α -D-Galactosamine:polypeptide N-acetylgalactosaminyltransferase that complements other GalNAc-transferases in complete O-glycosylation of the MUC1 tandem repeat. *J. Biol. Chem.* 273: 30472-30481.
- Online Mendelian Inheritance in Man, OMIM™. 1999. Johns Hopkins University, Baltimore, MD. MIM Number: 603565. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Hassan, H., et al. 2000. The lectin domain of UDP-N-acetyl-D-galactosamine: polypeptide N-acetylgalactosaminyltransferase-T4 directs its glycopeptide specificities. *J. Biol. Chem.* 275: 38197-38205.
- Sommer, P., et al. 2006. Identification of Tgf β 1i4 as a downstream target of Foxc1. *Dev. Growth Differ.* 48: 297-308.
- Wright, P.K., et al. 2009. Estrogen regulates vesicle trafficking gene expression in EFF-3, EFM-19 and MCF-7 breast cancer cells. *Int. J. Clin. Exp. Pathol.* 2: 463-475.
- O'Halloran, A.M., et al. 2009. Genetic polymorphisms in platelet-related proteins and coronary artery disease: investigation of candidate genes, including N-acetylgalactosaminyltransferase 4 (GALNT4) and sulphotransferase 1A1/2 (SULT1A1/2). *J. Thromb. Thrombolysis.* 27: 175-184.
- Lu, L., et al. 2009. GalNAc-transferase specificity prediction based on feature selection method. *Peptides* 30: 359-364.

CHROMOSOMAL LOCATION

Genetic locus: GALNT4 (human) mapping to 12q21.33.

SOURCE

GalNAc-T4 (N-12) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of GalNAc-T4 of human origin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-107552 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

GalNAc-T4 (N-12) is recommended for detection of GalNAc-T4 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for GalNAc-T4 siRNA (h): sc-96074, GalNAc-T4 shRNA Plasmid (h): sc-96074-SH and GalNAc-T4 shRNA (h) Lentiviral Particles: sc-96074-V.

Molecular Weight of GalNAc-T4: 66 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.